

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 December 2000 (28.12.2000)

PCT

(10) International Publication Number
WO 00/78794 A1

- (51) International Patent Classification⁷: C07K 5/08
- (21) International Application Number: PCT/US00/16864
- (22) International Filing Date: 19 June 2000 (19.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/338,409 22 June 1999 (22.06.1999) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 09/338,409 (CON)
Filed on 22 June 1999 (22.06.1999)
- (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).
- (72) Inventors; and
(75) Inventors/Applicants (for US only): MUNOZ, Benito [CA/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). SRINIVASAN, Kumar [IN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). WANG, Bowei [CN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).
- (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).
- (81) Designated States (national): CA, JP, US.
- (84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- Published:
— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 00/78794 A1

(54) Title: CYSTEINYL PROTEASE INHIBITORS

(57) Abstract: Amino acid di- and tri-peptide analogs having C-terminal vinylic groups, pharmaceutical compositions containing the analogs and methods of treating diseases using cysteinyl protease inhibitors; particularly neurodegenerative, autoimmune, cognitive disorders, stroke and traumatic injury.

CYSTEINYL PROTEASE INHIBITORS

FIELD OF THE INVENTION

The invention relates to protease inhibitor compounds useful in treatments of neurodegenerative, cardiovascular, pulmonary, neuromuscular, musculo-vascular and autoimmune diseases characterized by tissue damage, induction of apoptosis, cell death and/or the accumulation of amyloid proteins in deposits, plaques, fibrils and tangles.

BACKGROUND OF INVENTION

Cysteinyl proteases are implicated in important cellular processes including endocytosis and degradation of proteins, immune recognition, apoptosis and processing of amyloid precursor protein. Cysteinyl proteases are localized in the endosomal lysosomal compartment, possibly the endoplasmic reticulum and cytosol. While many of the molecular details are at present uncertain, the importance of cysteinyl proteases at the cellular level and at an organism level cannot be overstated.

In Alzheimer's disease (AD), amyloid precursor protein (APP; i.e., APP₆₉₅) is proteolytically processed in neural cells and A β cleavage fragments are secreted into cerebral extracellular perivascular matrix. Plaques containing A β act as a pathological landmark of AD (2-5) and *in vitro* studies have shown that A β can aggregate into insoluble fibrils which ultrastructurally resemble amyloid plaques (7). Aggregated in a fibrillar beta-pleated sheet configuration, A β is thought to form a nidus for binding other proteins contributing to Alzheimer's dementia and pathology. A β fibrils are reportedly toxic for cultured neurons *in vitro* (8), possibly through induction of

apoptosis (9), initiation of free radical formation (10) or increases in peroxide (H_2O_2) levels in cells (11).

APP, synthesized by neuronal cells, full length APP may be degraded in lysosomes (21,22), resulting in non-amyloidogenic fragments, or it can be proteolytically processed to polypeptides of 110kDa, i.e., sAPP (16-17); 9-10kDa (17); and 4.2kDa, i.e., $A\beta$ (18-19). Characterization of fragments and protease inhibitor profiles suggest three intracellular enzymes, termed secretases, differing as follows: namely,

Alpha-secretase (α -secretase) mediates cleavage of APP₆₉₅ and APP₇₅₁ producing sAPP (110kDa) and a 9-10kDa carboxy-terminal-fragment (CTF; i.e., APP₆₁₁₋₆₉₅). The CTF fragment may undergo further degradation in lysosomes (17). Products of lysosomal and α -secretase cleavages are generally considered to be non-amyloidogenic;

Beta-secretase (β -secretase) mediates cleavage of APP₆₉₅ in the carboxy-terminal region (i.e., at APP Met₅₉₅-Asp₅₉₆; $A\beta$ Asp₁; 23), producing a 11.4-11.8kDa fragment, 100 amino acids in length (CTF100), and containing the $A\beta$ domain; and,

Gamma-secretase (γ -secretase) mediates cleavage in the C-terminal transmembrane domain of APP (i.e., APP G₆₂₄AIIGLMVGGVVIA₆₃₇), and possibly also CTF100, to release the $A\beta_{1-39}$, $A\beta_{1-40}$ and/or $A\beta_{1-42}$ peptides. Both $A\beta_{1-40}$ and $A\beta_{1-42}$ are reportedly amyloidogenic; with $A\beta_{1-42}$ being more active than $A\beta_{40}$ (7,24,25).

Sensitivity of secretases to protease inhibitors has been studied in cell-based assays, i.e., by treating APP producer cells with protease inhibitors. Results have varied in different cells and under differing conditions of culture. In certain reports, APP cleavage is reportedly inhibited by serine protease inhibitors; in others by inhibitors of chymotrypsin-like enzymes; and in others by inhibitors of cysteinyl proteases such as cathepsins B, D, H, L and S and calpain. Studies with leupeptin (26,27), E64 and Z-Phe-Ala-CHN₂ (28), i.e., lysosomal enzyme inhibitors, have failed to implicate known lysosomal hydrolases in production of $A\beta$. Similarly, although calpain inhibitor I, a caspase inhibitor, was initially reported effective in

reducing A β levels *in vitro* (29), subsequent findings seem at odds with an effect on a putative γ -secretase (30). Dipeptidyl-aldehyde inhibitor, Z-L-Val-L/D-Phe-aldehyde (MDL 28170) and AEBSF [4-(2-aminoethyl)-benzenesulfonyl fluoride hydrochloride] were also once thought to inhibit A β_{1-42} production (32). Since non-specific acidotropic agents such as NH₄Cl and chloroquine, or Golgi inhibitors such as brefeldin A and monensin (19,27), reportedly affect APP processing it is possible that it takes place in the endoplasmic reticulum (ER) and/or Golgi apparatus. Specific and targeted inhibitors of secretase enzymes have (to date) proven elusive.

In apoptosis, an intracellular cascade of cysteinyl proteases, termed caspases, is activated by radiation damage to cellular DNA or structures (e.g., mitochondria), or by removal of one or more critical growth factors (e.g., EGF, PGF, TGF β , NGF, BDNF), glucose or serum. In addition, a variety of different receptor ligand interactions at the cell surface may trigger apoptosis *in vitro*: e.g., anti-Fas antibody or Fas ligand binding to Fas receptor; IL-1 binding to IL-1R β ; TNF α binding to TNFR1 and TNFR2; Apo 3 ligand binding to Death Receptor 3; glucocorticoids binding to cell surface glucocorticoid receptors; TRAIL, TRANCE and NOC18 binding to yet other cell surface "death" receptors. Important cysteinyl proteases activated in this manner are thought to include caspases 3, 8 and 9.

Apoptosis has recently been implicated in mechanisms by which tissue damage is triggered following traumatic injury, vascular insufficiency and stroke.

Using knockout mice, Cathepsin S is reportedly involved in class I MHC antigen processing, i.e., of self and viral antigens, while Cathepsin L is involved in class II processing of foreign proteins.




Thus, protease inhibitors find a variety of important medical and veterinary uses in treatments of diseases, as well as, uses in the food, dairy, agriculture, chemical and biotechnology industries.

SUMMARY OF THE INVENTION

Cysteinyll protease inhibitors, pharmaceutical compositions and methods of treating protein processing disorders, trauma, stroke, autoimmunity, neurodegenerative and cognitive disorders including Alzheimer's disease, are disclosed. The di- and tri-peptide analogues with unnatural α -amino side chain residues and derivatized amino and carboxyl terminal residues constitute highly effective inhibitors of cysteinyll proteases including e.g. cathepsins, caspases, calpains and secretases. Methods and pharmaceutical compositions are provided for modulating activities of cysteinyll proteases involved in diseases including methods for altering secretase-mediated processing of APP in Alzheimer's disease and aberrant cathepsin S-mediated processing of self proteins for MHC class I presentation in autoimmune diseases. The instant compounds include vinyl sulfides, sulfoxides, sulfones, amines, amine oxides, phosphines, phosphine oxides, phosphinates and phosphonates of peptidyl analogs of di- and tri-peptides, where the vinyl substituent in the peptidyl analog preferably occupies the carboxyl-terminal position and consists of a substituted alkenyl or alkynyl side chain.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Embodiments of the invention provide cysteinyll protease inhibitors. The instant compounds are vinyl heterocyclic alkyls and aryl adducts of peptidyl analogs of natural α -amino acid containing cysteinyll protease substrates; according to the following FORMULA I, II or III: namely,



wherein R_1 is an unnatural amino acid side chain, preferably a lower alkenyl or lower alkynyl that is unsubstituted, or alternatively, substituted with W,

W, in turn, is halo, hydroxy, alkyl, aryl, heterocycle, heteroaryl, alkoxy, aminosulfonyl, alkylcarbonyl, arylcarbonyl, heteroarylcarbonyl, heterocycle-carbonyl, nitro, haloalkyl; preferably, trifluoromethyl, cyano, amino or aminocarbonyl; and most preferably, a side chain not present in a natural amino acid;

X is $-\text{CH}=\text{CH}-(\text{CH}=\text{CH})_k-(\text{CH}_2)_j$ or $-\text{CH}_2\text{CH}_2-(\text{CH}_2)_b$ where "k" is 0-9, preferably 0-3, and most preferably 0; "j" is 0-4, preferably 0-2, most preferably 0; and, "b" is 0-6, preferably 3-6, most preferably 3 or 4;

Y is S, SO, SO_2 , NR_{20} , $-\text{N}(\text{O})(\text{R}_{20})-$, $-\text{PR}_{20}$, $-\text{P}(\text{O})(\text{R}_{20})-$, $-\text{P}(\text{O})\text{O}-$ or $-\text{P}(\text{O})(\text{OR}_{20})\text{O}-$; wherein, R_{20} is hydrogen, alkyl or aryl;

Z is $-(\text{CH}_2)_i-\text{A}$; "i" is 0-4, preferably 0-2, most preferably 0; wherein, A is a substituted or unsubstituted heterocycle, heteroaryl or aryl; where the substituents consist of one or more L groups; wherein, L is halogen, lower alkyl, lower alkenyl, lower alkynyl, alkoxy, aryloxy, hydroxy, haloalkyl, preferably trifluoromethyl, nitro, nitrile, alkylthio, phenyl, and $-\text{NR}_{30}\text{R}_{31}$; and, in optional embodiments L may be further substituted with W, as set forth above, but preferably L is selected from the substituents set forth in regard to V, below;

R_{30} and R_{31} are each independently H, alkyl, preferably lower alkyl, hydroxy or halo lower alkyl, and most preferably trifluoromethyl;

V is OH, halogen, lower alkyl, preferably methyl or ethyl or halogen-substituted lower alkyl, preferably halogen-substituted methyl or ethyl, and is more preferably OH;

R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , Q and n are each selected from (i), (ii), (iii), (iv), (v), (vi) and (vii) as follows;

(i) R_3 and R_5 are each independently selected from a side chain of a naturally occurring α -amino acid; H; alkyl, preferably lower (C_{1-6}) alkyl; alkenyl, preferably C_{2-10} alkenyl; alkynyl, preferably C_{2-6} alkynyl; aryl, aralkyl, aralkenyl, aralkynyl; heteroaryl, heteroaralkyl, heteroaralkenyl, heteroaralkynyl; L-substituted aryl, L-substituted aralkyl, L-substituted aralkenyl, L-substituted aralkynyl; M-substituted

heteroaryl, M-substituted heteroaralkyl, M-substituted heteroaralkenyl and M-substituted heteroaralkynyl; wherein,

M is lower alkyl, preferably C₁₋₄ alkyl, or alternatively, M is halo-lower alkyl, preferably C₁₋₄ haloalkyl, but most preferably M is trifluoromethyl;

R₂, R₄, R₆, and R₈ are each independently selected from among H; lower alkyl, preferably C₁₋₄ alkyl; cycloalkyl, preferably lower cycloalkyl; and cycloalkylalkyl, preferably lower cycloalkylalkyl;

R₇ is selected from among C₁₋₆ alkyl; R_AR_BCH; L-substituted or unsubstituted aryl; alkenyl; alkynyl; 9-fluorenyl; aralkyl; aralkenyl; aralkynyl; L-substituted or unsubstituted monocyclic or bicyclic heterocycle; L-substituted or unsubstituted monocyclic or bicyclic aryl, L-substituted or unsubstituted bicyclic aryl-heteroaryl, and L-substituted or unsubstituted monocyclic or bicyclic heteroaryl;

Q is -C(O)-, -O-C(O)-, -S(O)₂- or -HN-C(O)-;

n is zero or one;

R_A is -(T)_m-(D)_m-R₃, wherein T is O or NH, D is C₁₋₄ alkyl or C₂₋₄ alkenyl and m is zero or one; and

R_B is selected in the same manner as in regard to R₃ and R₅, above; or

(ii) R₂, R₅ and R₈ are selected as in (i), above;

R₃ and R₄ are selected as in either of (i), above, or (iv), below;

n is zero; and

R₆ and R₇ together with the atoms to which each is attached form a 3-10 membered ring, preferably a 4-8 membered ring, most preferably a substituted or unsubstituted 5-6 membered heterocyclic moiety containing 1-5 heteroatoms, where the substituents of the ring are selected either from those identified in regard to L, above, but, preferably those identified in regard to V, above, and most preferably the substituents so chosen will, together with the ring members will form a morpholino, thiomorpholino, pyrrolidinyl, V-substituted pyrrolidinyl, or 4-hydroxypyrrolidinyl ring; or

(iii) R₂, R₃, R₄, R₅ and R₈ are selected as set forth in regard to (i), above;

Q is -C(O)-

n is one; and

R₆ and R₇ are substituted or unsubstituted carbonyl (-(C = O)-), phenyl, a heteroatom, lower alkylene, preferably C₂₋₃ alkylene, or lower alkylene linked to a heteroatom, preferably C₂₋₃ alkylene linked to a heteroatom; together with the atoms to which they are attached R₆ and R₇ form a cyclic moiety, preferably a 4-6 membered cyclic or 8-12 membered bicyclic moiety, most preferably the subject moiety is succinimidyl, phthalimidyl or maleimidyl; and, wherein the substituents are selected from those set forth in regard to L, above, but preferably those set forth in regard to V, above, and

(iv) R₂, R₃, R₆, R₇, R₈, Q and n are selected as set forth above in regard to any of (i)-(iii), above, or (v)-(vii), below;

R₃ and R₄ are substituted or unsubstituted lower alkylene, preferably C₁₋₄ alkylene, or lower alkylene linked to a heteroatom, preferably C₂₋₄ alkylene linked to a heteroatom, or a heteroatom; together with the atoms to which they are attached, R₃ and R₄ form a heterocyclic moiety, preferably a 4-6 membered heterocyclic moiety, most preferably morpholino, thiomorpholino, pyrrolidinyl, or V-substituted pyrrolidinyl, particularly 4-hydroxypyrrolidinyl; wherein, the substituents of R₃ and/or R₄ are selected from those set forth in regard to L, above, but preferably to those set forth in regard to V, above; or

(v) R₂, R₇, R₈, Q and n are selected as set forth in regard to (i), above;

R₃ and R₄ are selected as in either of (i) or (iv), above;

R₅ and R₆ are each independently substituted or unsubstituted lower alkylene, preferably C₂₋₄ alkylene, or lower alkylene linked to a heteroatom, preferably C₂₋₄ alkylene linked to a heteroatom, or a heteroatom; together with the atoms to which they are attached R₅ and R₆ form a heterocyclic moiety, preferably a 4-6 membered heterocyclic moiety, most preferably a moiety selected from among morpholino, thiomorpholino, pyrrolidinyl, or V-substituted pyrrolidinyl, particularly 4-hydroxypyrrolidinyl; wherein, the substituents of R₅ and/or R₆ are selected from those set forth in regard to L, above, but preferably to those set forth in regard to V, above; or

(vi) R₂, R₆ and R₈ are selected as set forth in regard to (i), above;

R_3 and R_4 are selected as in either of (i) or (iv), above;

n is zero; and

R_5 and R_7 are substituted or unsubstituted lower alkylene, preferably C_{2-4} alkylene, or lower alkylene linked to a heteroatom, preferably C_{2-4} alkylene linked to a heteroatom, or a heteroatom; together with the atoms to which they are attached R_5 and R_7 form a heterocyclic moiety, preferably a 4-6 membered heterocyclic moiety, most preferably selected from among morpholino, thiomorpholino, pyrrolidinyl, or V-substituted pyrrolidinyl, particularly 4-hydroxypyrrolidinyl; wherein, the substituents of R_5 and/or R_7 are selected from those set forth in regard to L, above, but preferably to those set forth in regard to V, above; or

(vii) R_2 , R_3 and R_8 are selected as set forth in regard to (i), above;

R_3 and R_4 are selected as in either of (i) or (iv), above;

n is 0;

R_6 and R_7 together with the atoms to which each is attached form a 6-12, preferably an 8-10 membered bicyclic heterocyclic or heteroaryl moiety, preferably a reduced isoquinolinyl moiety, more preferably a 1,2,3,4-tetrahydroisoquinolinyl moiety;

and in all of (i)-(vii), above, unless specified otherwise: (a) the carbon chains, where so indicated, are straight or branched chain containing from 1 to about 12 carbons preferably 1 to about 6, and most preferably 4 to about 6 carbons; and (b) the cyclic moieties, where so indicated, contain one ring or two fused rings, each ring, or combination of rings, preferably containing 3 to about 16 atoms, most preferably 4 to about 12 atoms, and in a combination of two fused rings most preferably each of the subject rings contains 4 to about 6 ring atoms.

Abbreviations used are e.g., AD, Alzheimer's disease; A β , A β -amyloid peptide; APP, amyloid precursor protein; APP₇₅₁, wild-type human APP-751 protein; APP_{695WT}, wild-type human APP-695 protein; APP_{670/671}, human APP-695 protein harboring the "Swedish" double mutation at codons 670 and 671; APP_{670/671/717}, human APP-770 protein containing the "Swedish" double mutation at codons 670 and 671 and a Phe for Val substitution at codon 717 (codons numbered according to

the APP-770 amino acid isoform); α -sAPP, soluble APP; mAb, monoclonal antibody; PBS, phosphate-buffered saline; FBS, fetal bovine serum; SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis; DMSO, dimethylsulfoxide.

The terms used herein are intended to have meaning as follows: namely,

"Alkyl" is intended to mean straight, branched-chain or cyclic hydrocarbon chain composed of a number of carbon atoms that is commonly specified according to a formula. Where not specified, alkyl groups in the instant compositions preferably contain about 1 to about 12 carbon atoms (C_{1-12}). Representative examples of the subject alkyl groups include methyl, ethyl, propyl, cyclopropyl, isopropyl, n-butyl, t-butyl, sec-butyl, cyclobutyl, pentyl, cyclopentyl, n-hexyl, n-nonyl, n-decyl, cyclohexyl, cyclohexylmethyl, cyclohexylethyl, and the like.

"Alkenyl" is intended to mean a straight, branched-chain or cyclic hydrocarbon chain composed of one or more double-bonded carbon atoms, a number of which carbon atoms is commonly specified according to formula. Where not specified, alkenyl carbon chains in the instant compositions preferably contain about 2 to about 12 carbon atoms (C_{2-12}), and preferably about 2 to about 7 carbon atoms (C_{2-7}). The subject C_{2-12} alkenyl carbon chains contain about 1 to about 6 double bonds, while C_{2-7} alkenyl carbon chains preferably contain about 1 to about 3 double bonds. Representative alkenyl moieties include 2-methyl-2-propenyl, 2-methyl-1-propenyl, propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 2,2-difluoroethenyl, as well as, those straight and branched chain moieties having up to two double bonds. Cyclic alkenyl carbon moieties preferably contain one or two fused hydrocarbon ring moieties having about 3 to about 16 carbon atoms (C_{3-16}), and most preferably having about 4 to about 12 carbon atoms (C_{4-12}).

"Alkynyl" is intended to mean a straight, branched-chain or cyclic hydrocarbon chain composed of one or more triple-bonded carbon atoms, a number of which carbon atoms is commonly specified according to formula. Where not specified, alkynyl carbon chains in the instant compositions preferably contain about 2 to about 12 carbon atoms (C_{2-12}) with, preferably, about 1 to about 6 triple bonds, and the alkynyl carbon chains of 2 to 6 carbons (C_{2-6}), preferably, contain 1 to 3 triple

bonds. Representative alkynyl moieties include propynyl. Cyclic carbon alkynyl moieties preferably contain one or two fused hydrocarbon ring moieties having about 3 to about 16 carbon atoms (C_{3-16}), and most preferably about 4 to about 12 carbon atoms (C_{4-12}).

"Lower alkyl", "lower alkenyl" and "lower alkynyl" are intended to mean alkyl, alkenyl and alkynyl hydrocarbon chains, respectively (as defined supra), each of which chains contains fewer than six carbon atoms.

"Cycloalkyl" is intended to mean an unsaturated carbon bond single ring system moiety, or an unsaturated carbon bond multiple cyclic ring system moiety. Preferably, the subject cycloalkyl consists of about 3 to about 10 carbon atoms, and most preferably, the subject cycloalkyl consists of about 3 to about 6 carbon atoms. Representative example of the subject ring system moieties include one, two or more rings that are joined together in a fused, bridged or spiro-connected fashion, each of which moieties may be optionally substituted with one or more alkyl group substituents.

"Cycloalkenyl" is intended to mean a single ring system moiety, or a multiple ring system each of which moieties has at least one carbon-carbon double bond, and each of which moieties preferably contains about 3 to about 10 carbon atoms (C_{3-10}), and most preferably, about 4 to about 7 carbon atoms (C_{4-7}). Representative example of the subject ring system moieties include one, two or more rings that are joined together in a fused, bridged or spiro-connected fashion, each of which moieties may be optionally substituted with one or more alkyl group substituents.

"Cycloalkynyl" is intended to mean a single ring system moiety, or a multiple ring system each of which moieties has at least one carbon-carbon triple bond, and each of which moieties preferably contains about 3 to about 10 carbon atoms (C_{3-10}), and most preferably, about 8 to about 10 carbon atoms (C_{8-10}). Representative example of the subject ring system moieties include one, two or more rings that are joined together in a fused, bridged or spiro-connected fashion, each of which moieties may be optionally substituted with one or more alkyl group substituents.

"Heteroatoms" is intended to mean that the subject atoms are not the same. Where not specified, heteroatoms in the instant compositions are commonly non-

carbon atoms which are preferably are selected from among Oxygen (O), Nitrogen (N) and Sulfur (S). Substituents containing heteroatoms may be straight chain, branched chain or heterocyclic.

"Aryl" is intended to mean an aromatic hydrocarbon cyclic moiety, specified according to formula. Where not specified, in the instant compositions preferably the aryl moieties contain about 3 to about 14 carbon atoms (C_{3-14}) bonded into about 1 to about 2 ring structures. "Lower aryl" is intended to mean that the subject aryl moiety, defined supra, preferably contain about 5 to about 7 carbon atoms in the subject ring structure. "Aralkyl" is intended to mean that the subject aryl moiety, defined supra, is additionally substituted with one or more alkyl groups, as defined supra. Representative examples of aralkyl and aryl moieties include phenyl; benzyl; phenethyl; 1- and 2-naphthylmethyl; 1- and 2-naphthyl; 1- and 2-indenyl; pentalenyl; azulenyl; heptalenyl; acenaphthylenyl; 9-fluorenyl; phenalenyl; phenanthrenyl; anthracenyl; triphenylenyl; pyrenyl, chrysenyl; naphthacenyl; and the like.

"Heterocycle", and "heterocyclic", are intended to mean a cyclic carbon atom ring moiety composed of about 3 to about 14 atoms (C_{3-14}), preferably about 3 atoms to about 7 atoms constituting about 1 to about 2 ring moieties; of which atoms, one or more atoms are heteroatoms, as defined supra, and the subject preferred heteroatoms are as set forth above. Representative heterocycles include aliphatic saturated carbon ring moieties bonded with unsaturated heteroaryl ring moieties. Representative examples of heterocycles include pyrrolidiny, piperidiny, alkylpiperidiny, 1,4-dioxany, 1,4-dithianyl, 1,4-morpholinyl, 1,4-thiomorpholinyl, 1,2,3-oxadiazolyl, 1,3,5-trithianyl or 1,2,5-triazolyl.

"Heteroaryl" is intended to mean a cyclic carbon atom ring moiety composed of about 3 to about 14 atoms (C_{3-14}), preferably about 3 atoms to about 7 atoms constituting about 1 to about 2 ring moieties; of which atoms, one or more atoms is a heteroatom, as defined supra. The subject preferred heteroatoms are as set forth above, and most preferably at least one ring moiety comprises an aromatic ring. Representative examples of heteroaryls include furyl, thienyl, pyridyl, pyrrolyl, N-methylpyrrolyl, pyranyl, indolyl, quinoliny, isoquinoliny, cinnoliny,

quinoxaliny, naphthyridiny, benzothiazoly, dibenzothiazoly, xantheny and 1,2,3,4-tetrahydroisoquinoliny.

"NR^AR^B" is intended to mean a chemical moiety in which a nitrogen atom (N) is bonded to both an R^A substituent and an R^B substituent.

In certain alternative embodiments, one or more R group substituents, e.g., R₃, R₅, and the like, of Formula I, II or III, supra, may be identical with a side chain residue of a "naturally occurring amino acid". "Naturally occurring amino acids", and their constituent "side chain residues" are known in the art. "Natural side chain residues of naturally occurring amino acids", as used herein, is intended to mean natural substituents attached to the α -carbon atom of a naturally occurring amino acid synthesized by a plant or an animal. Representative examples of natural side chain residues include the hydrogen atom attached as a substituent at the α -carbon in glycine, the methyl group attached at the α -carbon in alanine, the isopropyl group attached at the α -carbon in valine, and the like.

"Halogen", "halide" and/or "halo" are used interchangeably to refer substituents selected from among a fluorine atom (F), chlorine atom (Cl), bromine atom (Br), iodine atom (I), as well as pseudohalide atoms. Where not specified, halogen in the instant compositions preferably contain one or more atoms selected from among fluorine, chlorine, bromine and iodine.

"Pseudohalides" is intended to mean atoms, or chemical constituent groups, that have one or more chemical properties and/or chemical reactivities substantially similar to a halogen. The subject constituent groups are distinguished by being interchangeable with halogens in the instant synthetic methods, as disclosed further below. Representative examples of pseudohalides include, but are not limited to, cyanide, cyanate, thiocyanate, selenocyanate, azide and trifluoromethyl.

"Haloalkyl" is intended to mean an alkyl group substituted with one or more halogens.

"Halo-lower alkyl" is intended to mean an alkyl moiety containing six or fewer carbon atoms which is substituted with one or more halogens. Where not

explicitly specified otherwise, preferred halo-lower alkyl moieties in the instant compositions are selected from among $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, $-\text{CH}_2\text{Cl}$ and $-\text{CH}_2\text{Br}$.

"Haloalkoxy" is intended to mean an $\text{RO}-$ moiety in which R is a haloalkyl group.

"Aminocarbonyl" is intended to mean an $-\text{C}(\text{O})\text{NH}_2$ moiety.

"Alkylaminocarbonyl" is intended to mean an $-\text{C}(\text{O})\text{NHR}$ moiety in which R is hydrogen, alkyl, preferably lower alkyl or aryl, preferably lower aryl.

"Dialkylaminocarbonyl" is intended to mean an $-\text{C}(\text{O})\text{NR}'\text{R}$ moiety in which R' and R are independently alkyl or aryl, preferably lower alkyl or lower aryl;

"Carboxamide" is intended to mean groups of formula $-\text{NR}'\text{COR}$.

"Alkoxy" is intended to mean an $-\text{C}(\text{O})\text{OR}$ group in which R is alkyl, preferably lower alkyl or aryl, preferably lower aryl.

"Alkoxy" is intended to mean an $\text{RO}-$ group in which R is alkyl, preferably a lower alkyl or aryl, and most preferably a lower aryl.

"Thioalkoxy" is intended to mean an $\text{RSO}-$ group in which R is alkyl, preferably a lower alkyl or aryl, and most preferably a lower aryl.

"Alkylene" is intended to mean a straight, branched or cyclic hydrocarbon moiety, preferably a straight or branched moiety composed of about 1 to about 20 carbon atoms (C_{1-20}), and most preferably composed of fewer than about 12 carbon atoms (C_{1-12}). In presently preferred embodiments, the alkylene moiety is either unsubstituted, or alternatively, is substituted with one or more alkyl groups, as defined supra. In certain alternative embodiments of the invention, the subject hydrocarbon moiety making up an alkylene additionally contains one or more heteroatoms selected from among oxygen and nitrogen. In other embodiments, the latter subject nitrogen heteroatom can constitute an amide nitrogen bonded with an alkyl group. Representative examples of alkylene groups include methylene ($-\text{CH}_2-$), ethylene ($-\text{CH}_2\text{CH}_2-$), propylene ($-(\text{CH}_2)_3-$), cyclohexylene ($-(\text{C}_6\text{H}_{10})-$), methylenedioxy ($-\text{O}-\text{CH}_2-\text{O}-$) and ethylenedioxy ($-\text{O}-(\text{CH}_2)_2-\text{O}-$).

"Lower alkylene" is intended to mean an alkylene moiety composed of fewer than about 6 carbon atoms. In certain preferred embodiments, lower alkylene groups contain about 2 to about 3 carbon atoms.

"Alkenylene" is intended to mean a straight, branched or cyclic hydrocarbon, preferably straight or branched chain composed of about 2 to about 20 carbon atoms (C_{2-20}) at least two of which are double bonded one to another. In certain presently preferred embodiments of the invention, the subject alkenylene moiety is preferably composed of about 2 to about 12 carbon atoms (C_{2-12}), and in most preferred embodiments the subject moiety preferably consists of a "lower alkenylene" having about 2 to about 6 carbon atoms. In certain alternative embodiments, the subject alkenylene moiety may be composed of a compound substituted at one or more positions in the straight, branched or cyclic hydrocarbon with one or more alkyl groups, as defined supra. In other alternative embodiments, the subject straight, branched or cyclic hydrocarbon may contain one or more heteroatoms, i.e., in place of one or more carbon atoms. In the latter case, the subject heteroatoms are preferably oxygen, sulfur or nitrogen atoms. The latter nitrogen atom may, in other alternative embodiments, constitute an amide nitrogen bonded with an alkyl group. Representative examples of alkenylene groups include -CH=CH-CH=CH-, -CH=CH- and -CH=CH-CH₂-.

"Lower alkenylene" is intended to mean an alkenylene having about 2 to about 6 carbon atoms, and preferably about 2 to about 4 carbon atoms.

"Alkynylene" is intended to mean a straight, branched or cyclic hydrocarbon, preferably straight or branched chain composed of about 2 to about 20 carbon atoms (C_{2-20}) at least two of which are triple bonded one to another. In certain presently preferred embodiments, the subject alkynylene moiety preferably comprises about 2 to about 12 carbon atoms (C_{2-12}), and in most preferred embodiments the subject moiety consists of a "lower alkynylene" having 2 to about 6 carbon atoms. In certain alternative embodiments, the subject alkynylene moiety may be composed of a compound substituted at one or more positions in the straight, branched or cyclic hydrocarbon with one or more alkyl groups. In other alternative embodiments, the subject straight, branched or cyclic hydrocarbon may contain one or more heteroatoms, i.e., in place of one or more carbon atoms. The latter heteroatoms are preferably oxygen, sulfur or nitrogen atoms. The latter nitrogen may, in other alternative embodiments, constitute an amide nitrogen bonded with an "alkyl group

substituent", as defined below. Representative examples of alkynylene groups include $-C\equiv C-C\equiv C-$, $-C\equiv C-$ and $-C\equiv C-CH_2-$.

"Lower alkynylene" is intended to mean an alkynylene composed of about 2 to about 6 carbons. In presently preferably embodiments of the invention, lower alkynylene moieties are composed of about 2 to about 4 carbon atoms.

"Arylene" is intended to mean a monocyclic or polycyclic hydrocarbon ring moiety having one or more aromatic/aryl groups as constituents in the subject ring moiety, and additionally, having one or more carbon-carbon double bonds. The arylene moiety is preferably composed of 3 to about 20 carbon atoms (C_{3-20}). In presently most preferred embodiments, the subject arylene groups are composed of about 3 to about 12 carbon atoms (C_{3-12}), and most preferably, the arylene group is a "lower arylene" (C_{5-6}). In alternative embodiments, the subject arylene moiety may, in turn, be substituted with one or more aryl groups, and the arylene cyclic ring form may also, optionally, contain one or more heteroatoms. The latter heteroatoms constituting the arylene cyclic ring are preferably oxygen, sulfur or nitrogen, and the subject nitrogen atom may, in other embodiments, constitute an amide nitrogen bonded to an "alkyl group substituent". Representative examples of arylene groups include 1,2-, 1,3- and 1,4-phenylenediamine.

"Lower arylene" is intended to mean an arylene group having about 8 to about 10 carbon atoms (C_{8-10}).

"Alkylidene" is intended to mean a $=CR'R''$ substituent group, in this case, attached to a basal hydrocarbon core structure such as a $=CR'R''$ group substituent of an alkylene or a heterocycleene. In the latter case, the $=CR'R''$ substituent group is attached to the core structure through the group's carbon-carbon double bond ($=C-$). Representative examples of alkylidene groups include methylenide ($=CH_2$) and ethylenide ($=CHCH_3$).

"Arylalkylidene" is intended to mean an alkylidene group, i.e., $=CR'R''$, in which either the R' or the R'' substituent constitutes an aryl group.

"Carbonyl" is intended to mean a bivalent carbonyl group of the general formula $-(C=O)-$.

"Alkyl group substituent" is intended to mean a hydrocarbon, a halo-hydrocarbon or a halogen-containing substituent to a core structure as follows: namely, in presently preferred embodiments, the subject "alkyl group substituent" is halo-lower-alkyl, aryl, hydroxy, alkoxy, aryloxy, alkylthio, arylthio, aralkyloxy, aralkylthio, carboxy, alkoxycarbonyl, an oxygen atom or a cycloalkyl group.

"Aryl group substituent", when used in regard to a core structure that is alkyl, cycloalkyl, cycloalkylalkyl, aryl, or a heteroaryl moiety, is intended to mean that the subject core structure is substituted with 1 or more, and preferably 1 to about 3, "aryl group substituents" selected from the following: namely, halogen atoms, alkyls, halo-alkyls, arylalkyl, heteroarylalkyl, alkenylene, alkynylene and hydroxyl. In presently preferred embodiments of the invention wherein the core structure is a halo-lower-alkyl, the preferred "aryl group substituents" are trifluoromethyl, formyl, alkylcarbonyl or substituted or unsubstituted arylcarbonyl. In the latter case the substituents comprise 1 or more, and preferably 1 to about 3 groups selected from among the following: namely, halogen atoms, halo-alkyls, alkyls, heteroarylcarbonyls, carboxyls; alkoxycarbonyls, aryloxy carbonyls, aminocarbonyls, alkylaminocarbonyls, dialkylaminocarbonyls, arylamino-carbonyls, diarylaminocarbonyls, arylalkylaminocarbonyls, alkoxys, aryloxys, perfluoroalkoxys, alkenyloxys, alkynyloxys, arylalkoxys, aminoalkyls, alkylaminoalkyls, dialkylaminoalkyls, arylaminoalkyls, amino groups, alkylamino groups, dialkylamino groups, arylaminos, alkylarylaminos, alkylcarbonylaminos, arylcarbonyls-, amines, azidos, nitrogen atoms, thiol groups, alkylthiol groups, arylthiols, perfluoroalkylthiols, thiocyano groups, isothiocyano groups, alkylsulfinyl groups, alkylsulfonyls, arylsulfinyls, arylsulfonyls, aminosulfonyls, alkylaminosulfonyls, dialkylaminosulfonyls and arylaminosulfonyl groups.

"Aminosulfonyl" is intended to mean either an $\text{-NHSO}_2\text{-}$ substituent or an $\text{-SO}_2\text{NH-}$ bivalent group.

"Amido" is intended to mean either a -C(O)NH- or an -HNC(O)- bivalent group.

"Thioamido" is intended to mean either a -C(S)NH- or an -HNC(S)- bivalent group.

"Oxyamido" is intended to mean a -OC(O)NH- or an -HNC(O)O- bivalent group.

"Thiaamido" is intended to mean a -SC(O)NH- or an -HNC(O)S- bivalent group.

"Dithiaamido" is intended to mean a -SC(S)NH- or an -HNC(S)S- bivalent group.

"Ureido" is intended to mean an -HNCONH- bivalent group.

"Thioureido" is intended to mean an -HNCSNH- bivalent group.

"Alzheimer's disease" is the most common form of dementia affecting nearly 25% of the elderly population over the age of 85 years. Alzheimer's disease (AD) can now be diagnosed with a considerable degree of accuracy with 90% correspondence between clinical diagnosis and autopsy confirmation at certain centers. The characteristic histopathologic changes at autopsy include neurofibrillary tangles, neuritic "senile" plaques, neuronal loss and amyloid angiopathy (1). The major constituent of plaques (plaques) is a 4.2kDa fibrillogenic polypeptide of 42-43 amino acids in length and referred to as amyloid beta-protein (abbreviated A β). A β is derived by proteolytic cleavage from amyloid precursor protein (APP; 2-5) and accumulates in the cerebral extracellular perivascular matrix in patients with AD, where its observance is considered to be a pathological landmark of the disease. When aggregated in fibrillar beta-pleated sheet configurations, A β is thought to form a nidus for binding of other proteins that may contribute to pathology. Binding of A β to neurons can result in increased vulnerability to excitotoxicity, generation of oxygen free radicals, impaired membrane transport systems, inhibition of glutamate receptors and induction of apoptosis (6). Neuronal loss can lead to disruption and defects in neurotransmission, cell-cell communications and abnormal production of extracellular matrix proteins. In addition, deposition of A β in and around cerebral blood vessels, i.e., cerebral amyloid angiopathy, has been associated with hemorrhagic stroke in the elderly and patients with AD.

"Amyloid precursor protein", abbreviated APP, is intended to mean an amino acid sequence of Accession No. 1070623 (Entrez: www.ncbi.nlm.nih.gov), having

695 or 770 amino acids, and sequences related thereto e.g. by conservative amino acid substitution and the like. Representative proteins have one or more of the following attributes: namely, a calculated theoretical molecular weight about 86,943, encoded by a gene mapping to 21q21.2 and 2-21q21.2, alternatively spliced to encode either a protein of 695 aa or 770 aa as a C-terminal extended polypeptide with a Kunitz-type serine protease inhibitor domain, Gly₇₀₀-Leu₇₂₃ being the transmembrane domain, and Lys₇₂₃-Phe₆₉₅ or Lys₇₂₃-Asn₇₇₀ being the cytoplasmic domain.

"A β domain", also referred to as A β -peptide", and abbreviated "A β ", is intended to mean a cleavage product of APP, supra, as mediated by one or more proteases at an APP NH₂-terminal Lys₆₇₀-Met₆₇₁ or Met₆₇₁-Asp₆₇₂ residue and a COOH-terminal Val₇₁₀-Val₇₁₁ or at a COOH-terminal Ala₇₁₃-Thr₇₁₄. "A β ₁₋₄₀" being Asp₆₇₂-Ile₇₁₂; "A β ₁₋₃₉" being Asp₆₇₂-Val₇₁₁; A β ₁₋₄₂ being Asp₆₇₂-Thr₇₁₄; and "A β ₁₋₄₃" being Asp₆₇₂-Val₇₁₅.

"Carboxy terminal fragment", abbreviated interchangeably CTF and i β -CTF, is intended to mean the portion of APP (Accession No. 1070623) corresponding to an APP fragment having a molecular size on SDS-PAGE of about 9kDa to about 12kDa and/or a sequence from about Asp₆₇₂ to about Asn₇₇₀.

"CTF 100", meaning carboxy terminal fragment of 100 amino acids in length, is intended to mean an APP fragment having an apparent molecular size on SDS-PAGE of about 9kDa to about 12kDa and a sequence from about Asp₆₇₂ to about Asn₇₇₀.

"Soluble APP", abbreviated sAPP, is intended to mean the portion of APP (Accession No. 1070623) corresponding to an N-terminal fragment produced by α -secretase having a molecular size on SDS-PAGE of about 100 kDa to about 115 kDa and lacking the transmembrane (supra) and cytoplasmic (supra) domains of APP.

"Treatment" is intended to mean a method of delivering to a subject in need thereof a pharmaceutical preparation according to the compounds of the invention with the aim of effecting a change in APP:A β processing in the subject in need thereof. The instant methods include delivering the preparation to a patient i) before the dysfunction has been diagnosed, e.g., prophylactic protocols delivered with the

aim of preventing development of the dysfunction, as well as, ii) after the dysfunction has been diagnosed, e.g., therapeutic protocols. That the subject treatments have fulfilled the intended aim of effecting a change in APP processing to A β in the subject will be evident by one or more of the following: namely, (i) a change in the amount of A β or sAPP in a sample of a biological tissue or biological fluid collected from a treated patient; (ii) a change in the amount of one or more APP:A β processing proteins in a biological tissue or biological fluid collected from a treated patient; or, (iii) a change (increase or decrease) or complete elimination of one or more clinical indicia of disease, e.g., diagnostic symptoms, in a treated patient.

"Dysfunction" is intended to mean a pathologic condition of APP processing to A β in a tissue, i.e., one or more changes in APP processing and/or production of A β as compared with the processing of APP occurring in a normal healthy control cohort. For example the subject pathological conditions include, but are not limited to, i) toxic amyloid dystrophy, (e.g., chemical or drug-induced secondary amyloid dystrophy), ii) vascular impairment with amyloid deposition, iii) degeneration and peripheral degeneration of vascular and/or neural tissue associated with amyloid deposition, iv) cellular detachment from basement membrane and defects in tissue organization visible microscopically in histology sections, e.g., in the endothelium of vessels; v) amyloid containing lesions resulting following physical trauma; and v) hereditary amyloid deposition in tissues, e.g., systemic amyloidosis and familial Alzheimer's diseases.

"Subject in need thereof" is intended to mean a mammal, e.g., humans, domestic animals and livestock, having one or more dysfunctions of APP processing, as defined above.

"Biological fluid" as used herein is intended to mean a tissue fluid, such as cerebrospinal fluid (CSF); blood, plasma and serum; fluid collected from a body cavity (i.e., peritoneal fluid, lung lavage fluid, urogenital mucus secretions, and the like); urine, feces, sputum, sweat; and the like.

"Processing of APP to A β ", abbreviated APP:A β processing, is intended to mean intracellular pathways by which proteases act upon APP to catalyze hydrolysis

of peptide bonds thereby liberating the A β fragment (as defined supra) of APP; in this case, including processes active in the following intracellular compartments: namely, in the endoplasmic reticulum; in the cytoplasm associated with ribosomes; in cytoplasmic vesicles (e.g., Golgi secretory vesicles); and, in lysosomal vesicles and the like.

"Substantially pure" is intended to mean that the subject preparation is sufficiently homogeneous to appear free of detectable impurities as determined by standard methods of physicochemical analysis, e.g. IR, NMR, spectral methods for determining chirality and the like, or by separation methods, e.g., thin layer chromatography (TLC), high performance liquid chromatography (HPLC), gas liquid chromatography (GLC) such as subject to analysis by mass spectrometry (GCMS), or by another method employed to assess chemical purity. A substantially chemically pure compound may, however, be a mixture of stereoisomers. In such instances, further purification may increase the specific activity of the compound e.g. by isolating a single isostere.

"Prodrug" is intended to mean one or more of the instant compounds in a form that, upon *in vivo* administration, is metabolized and/or otherwise converted from a form lacking in a biological or pharmaceutical activity, into a form having the subject activity. Representative examples of forms of prodrugs are disclosed e.g. in Nogrady (1985) *Medicinal Chemistry A Biochemical Approach*, Oxford University Press, New York, at pages 388-392.

"Isostereoisomer", abbreviated isostere, is intended to mean of a compound possessing of one of two or more stereochemical attributes expressed by one of the instant compounds, e.g. one spectral property. An isostere compound may exhibit a biologic and/or a pharmacologic activity that is greater than, or less than that of a racemic mixture of one of the instant compounds.


"Alter APP processing to A β ", is intended to mean altering the processing of APP. Two alternative processing pathways for APP may be represented as follows (the arrows represent cleavage sites):

ingredient in the composition or may be combined with other active ingredients. Liposome delivery systems are also contemplated, e.g., as disclosed in U.S. Patent No. 4,522,811.

The concentration of active compound(s) in the instant pharmaceutical composition will depend on absorption, inactivation and excretion rates of the active compound(s) (i.e., the selected compounds of FORMULA I, II and/or III, above), the physicochemical characteristics of the compound(s), the dosage schedule, and amount administered as well as other factors known to skilled artisans. Typically a therapeutically effective dosage will produce a serum concentration of one or more of the instant compounds ranging from about 0.1 ng/ml to about 50-100 µg/ml serum or plasma. The pharmaceutical compositions typically should provide a dosage of from about 0.001 mg to about 2000 mg of compound per kilogram (kg) of body weight per day. Pharmaceutical dosage unit forms may be prepared to provide about 1 mg to about 1000 mg, preferably about 10 mg to about 500 mg of the instant compound. The dosage and method of treatment will be determined by the physician after consideration of the general health, size and condition of a patient.

Effective concentrations (or amounts) of one or more of the compounds described herein or pharmaceutically acceptable derivatives thereof may be mixed with one or more suitable pharmaceutical carriers or vehicles. Solutions or suspensions useful for parenteral, intradermal, subcutaneous, or topical administration may include a sterile diluent such as water for injection, saline solution; a fixed oil; a polyethylene glycol; a glycerin; a propylene glycol or other synthetic solvent; one or more antimicrobial agents, e.g. benzyl alcohol and methyl parabens; an antioxidant, e.g. ascorbic acid or sodium bisulfite; a chelating agent, e.g. ethylenediaminetetraacetic acid (EDTA); a buffer, e.g. an acetate, citrate or phosphate buffer; or, an isotonic agent e.g., sodium chloride or dextrose. Parenteral preparations may be enclosed in ampules, disposable syringes or single or multiple dose vials made of glass, plastic or other suitable materials.

Methods for improving the solubility of the instant compounds, e.g., during administration and delivery from within the instant pharmaceutical compositions, include but are not limited to, uses of co-solvents, e.g., dimethylsulfoxide (DMSO).






"Altering processing of APP to A β ", abbreviated altering APP:A β processing, when used in regard to a composition containing one or more of the instant compounds is intended to mean that the subject composition is effective when administered to a cell, e.g., in tissue culture, to inhibit the production of carboxy terminal fragments of APP including i β -CTF and/or A β . While not wishing to be tied to any particular mechanism of action, representative examples of mechanisms by which the latter inhibitory effect on A β production may be exerted by the instant compounds include inhibiting: (a) an enzyme activity in a cell, e.g., an enzyme exhibiting a γ -secretase activity; (b) a co-factor binding to, and exerting an effect on, the subject enzyme activity in the cell, e.g., a cofactor increasing the catalytic efficiency or turnover rate of an enzyme with a γ -secretase activity in a cell; (c) a transcription regulator, e.g., altering the level of expression of mRNA encoding the subject enzyme or cofactor protein in a cell; (d) a protein kinase, e.g., altering either the level of expression of, or the activity of, the subject enzyme in a cell; and (e) a phosphorylase, e.g., altering the activity of the subject enzyme in the cell. The subject compositions may act directly on enzyme:cofactor:APP substrate complexes, or they may act indirectly, i.e., to change transcription or translation of enzyme proteins or cofactors.

"Altering apoptosis" is intended to mean that the subject compound is effective to inhibit a cysteinyl protease activated in a cell which has been treated to induce programmed cell death. Representative proteases so activated include calpains, caspases (e.g., caspases 3, 8 and 9), cathepsins and the like. Representative apoptosis inducer agents so able to induce apoptosis include Fas antibody, staurosporine, interleukins IL-1 and TNF, glucose and oxygen deprivation during ischemic tissue injury and the like. Apoptosis may be induced in the cells of tissues subject to traumatic injury, restricted blood flow (e.g., during peripheral microvascular clotting and stroke) and the like.

"Apoptosis" is intended to mean the cascade of energy (ATP) dependent events triggered by an apoptosis inducer agent and leading to programmed cell death through mechanisms commonly involving intracellular caspase enzymes, and commonly requiring about 12 to about 24 hours to accomplish the subject cell death. Apoptosis does not involve rapid, non-energy dependent lysis of cells, such as occurs during necrosis with release of cellular constituents (e.g., LDH) into the extracellular space. However, during apoptosis cells fragment their DNA and package it into membrane vesicles.

Embodiments of the invention provide presently preferred compounds of FORMULA I, II and III as set forth according to FORMULAS IV, V and VI: as follows,

namely,



wherein, all of the groups and substituents thereof, e.g. R_1 - R_8 , Q, Y and Z are identical with those set forth in regard to FORMULA I, II and III, above. In certain alternative preferred embodiments of compounds according to FORMULA IV, V or VI, when $(\text{Q})_n$ is a carbonyl group and R_1 and R_7 and the atoms to which they are attached form a heterocyclic ring moiety, the presently preferred ring moieties are succinimide, phthalimide or maleimide, and the most preferred is phthalimide. Similarly, when n is zero and R_1 , R_7 and the atoms to which they are attached form a heterocyclic ring moiety, the preferred heterocyclic ring moieties are morpholino, thiomorpholino, pyrrolidinyl and V-substituted pyrrolidinyl, and most preferably

4-hydroxypyrrolidinyl and 1,2,3,4-tetrahydroisoquinolinyl. When n is zero and R_3 , R_1 or R_2 , and R_7 taken together with the atoms to which they are attached form heterocyclic moieties, then the preferred ring moieties are morpholino, thiomorpholino, pyrrolidinyl, and V -substituted pyrrolidinyl, and most preferred is 4-hydroxypyrrolidinyl. Representative examples of the presently preferred compounds include: e.g., N-Cbz-L-Leu-L-Leu-L-Leuene-phenyl vinyl sulfone; N-Cbz-L-Leu-L-Leu-D-Leuene-phenyl vinyl sulfone; N-(THIQ-carbonyl)-L-Leu-L-Leu-D-Leuene phenyl vinyl sulfone; N-(THIQ-carbonyl)-L-Leu-L-Leu-L-Leuene phenyl vinyl sulfone, N-(THIQ-carbonyl)-L-Val-L-Met-L-Leuene phenyl vinyl sulfone; N-(THIQ-carbonyl)-L-Val-L-Met-D-Leuene phenyl vinyl sulfone; N-(THIQ-carbonyl)-L-Val-L-Leu-L-Leuene phenyl vinyl sulfone; N-(THIQ-carbonyl)-L-Val-L-Leu-D-Leuene phenyl vinyl sulfone; N-(4-benzylpiperidinyl-carbonyl)-L-Leu-L-Leu-L-Leuene phenyl vinyl sulfone; N-(4-benzylpiperidinyl-carbonyl)-L-Leu-L-Leu-D-Leuene phenyl vinyl sulfone; N-(4-benzylpiperazinyl-carbonyl)-L-Leu-L-Leu-L-Leuene phenyl vinyl sulfone; and, N-(4-benzylpiperazinyl-carbonyl)-L-Leu-L-Leu-D-Leuene phenyl vinyl sulfone.

Embodiments of the invention provide pharmaceutical compositions in which the instant compounds are present as the salts of organic acids and bases (such as acetates, lactates, tartrates, citrates, ascorbates, succinates, butyrates, valerates, fumarates and the like); salts of inorganic acids and bases, e.g., alkali metal salts of lithium, potassium, sodium, alkaline earth metals, such as barium, calcium and magnesium and the like, salts of transition metals such as Zinc, salts of mineral acids such as hydrochlorides and sulfates; esters; acids; bases; solvates; hydrates; amine salts; phosphonates such as sodium hydrogen phosphate and disodium phosphate, and the like; including hydrolyzable prodrugs cleavable and activated *in vivo* to drugs, e.g., derivatives of N,N' -dibenzylethylenediamine, chlorprocaine, choline, ammonia, diethanolamine and other hydroxyalkylamines, ethylenediamine, N -methylglucamine, procaine, N -benzylphenethylamine, 1-para-chlorobenzyl-2-pyrrolidin-1'-yl-methylbenzimidazole, diethylamine and other alkylamines, piperazine and tris(hydroxymethyl)aminomethane and the like.

In other embodiments the invention provides a pharmaceutical preparation consisting of one or more isolated and substantially pure diastereoisomers or chiral forms of a pharmaceutical preparations according to FORMULA (I), (II) and/or (III).

Embodiments of the invention provide compounds according to FORMULA I, II or III that are effective to modulate the activity of cysteinyl proteases including extracellular (secreted) proteases and intracellular protease involved in protein processing. The instant protease inhibitors include inhibitors acting, e.g., by processes including both competitive and non-competitive inhibition.

In yet other embodiments, the invention provides methods for administering to a subject in need thereof a concentration of one or more pharmaceutical preparations consisting of one or more compound according to FORMULA I, II and/or III that is effective to ameliorate one or more symptoms of disease. Examples of subjects so-in-need include e.g. patients with systemic amyloidosis, Alzheimer's disease, atherosclerosis, restenosis following angioplasty, rheumatoid arthritis and connective tissue diseases, dermatologic diseases such as Epidermolysis bullosis, Systemic lupus erythematosus, Sjogren's syndrome, Insulin-dependent diabetes types I and II, atherosclerosis, stroke, traumatic neural degeneration, chronic vascular and respiratory disorders, necrotic and apoptotic tissue damage resulting from bacterial, fungal and viral infection, (e.g., AIDS dementia, adenovirus, HSV, HPV, influenza and the like), inflammatory bowel disease, irritable bowel syndrome, diarrhea, constipation, gastric acid secretion and ulcers, complications resulting from co-medication or surgical procedures; or heart insufficiency, hyperprolactinemia, asthma and arthritis.

Embodiments also provide compounds according to FORMULA I, II or III. that are capable of inhibiting the activity of a cysteinyl protease, e.g., an intracellular secretase such as β -secretase involved in processing of APP and production of A β , an intracellular apopain/caspase protease involved in programmed cell death (apoptosis), an intracellular protease involved in antigen processing, e.g. a cathepsin, or an extracellular protease involved in tissue destructive processes, e.g., in trauma or arthritis. That a compound is so active may be determined by conducting e.g. either an *in vitro* or an *in vivo* assay. For example, in an *in vitro* assay the instant

compounds decrease in the activity of a cellular cysteinyl protease enzyme activity, or a secreted extracellular cysteinyl protease. Assays for cysteinyl proteases, as with assays for other proteases, are within the ordinary skill in the art. In one representative example, the subject cysteinyl protease is an intracellular β -secretase enzyme involved in processing of APP and production of $A\beta$ amyloidogenic polypeptides. In the latter case a positive assay is recorded when one or more of the following are observed: namely, (1) a decrease is observed in the 4-kDa amyloid $A\beta$ in the medium relative to control cultures; and/or (2) the relative amount of sAPP, in the medium is observed to increase; (3) a decrease is observed in the amount of C-terminal APP fragments >9 kDa in cell lysates, i.e., resulting from differential processing; and/or (4) there is an observed increase in the amount of α -sAPP in the medium relative to control cultures. *In vivo*, the instant compounds are preferably effective when administered to an experimental animal of AD or systemic amyloidosis to either: (1) increase the levels of sAPP (e.g., in brain, plasma or CSF), and/or (2) decrease the level $A\beta$ in a biological fluid or tissue extract. In one representative example, a positive assay result may be indicated as an increase in the ratio of α -sAPP to sAPP in a tissue sample from a pre-treatment to a post-treatment interval.

Representative methods for preparing the instant vinyl sulfones are depicted in the EXAMPLES, section below, and in particular in Reaction Schemes A-F of EXAMPLES 1-14. In Scheme A, preparation of amino ester hydrochlorides is depicted; Scheme B illustrates preparation of vinyl sulfones from the amino ester hydrochlorides; in Scheme C, formation of di- and tri-peptide vinyl sulfone analogs is depicted; Scheme D illustrates the preparation of an illustrative isomer of an amino vinyl sulfone; and, synthesis of illustrative N-protected tripeptide vinyl sulfone analogs as 1,2,3,4-tetrahydro-isoquinolinyl ureas is shown in Scheme E. An alternative to the synthesis of Scheme E is presented in Scheme F, i.e., 4-benzylpiperidine or 1-benzylpiperazine are substituted for 1,2,3,4,5-tetrahydroisoquinoline. As an alternative, an automated peptide synthesizer and solid phase sequential coupling procedures may be used to produce intermediates useful in

final synthesis of the instant di- and tri-peptide inhibitors. For example, the instant compounds may be synthesized from their carboxyl terminal end toward their amino terminal end using sequentially protected amino acids, amino acid derivatives and/or isomers, or spacer groups, according to the structure set forth in FORMULAS I, II and/or III. A variety of solid phase resin supports are useful in this method including certain of those useful in the art for solid phase synthesis of peptides. One presently preferred resin comprises polystyrene cross-linked with from about 0.5% to about 3% modified divinyl benzene, wherein the modification involves benzhydrylamidation, chloromethylation or hydroxymethylation. Modification is effective to produce sites that are useful for linking the instant di- and tri-peptide intermediates, e.g., through amide or ester bonds. One representative example of a hydroxymethyl resin is disclosed by Bodansky *et al.* W 966) *Chem. Ind.* (London) 38:1597-98). Preparation of illustrative chloromethyl- and benzhydrylamine-resins is disclosed in Stewart *et al.* ("*Solid Phase Peptide Synthesis*" 2nd Edition, Pierce Chemical Co., Rockford, Illinois (1984), Chapter 2, pp. 54-55). Illustrative methods for coupling protected amino acids to a resin are disclosed by Gisin ((1973) *Helv. Chem. Acta* 56:1476).

A variety of α -amino protecting groups may be useful in synthesis of the instant compounds and their intermediates. Representative examples of such protecting groups as presently contemplated include: (i) acyl protecting groups such as formyl, trifluoroacetyl, benzoyl, phthalyl, p-toluenesulfonyl (tosyl), benzenesulfonyl, nitrophenylsulfonyl, tritylsulfonyl, o-nitrophenoxycarbonyl, α -chlorobutyryl, 4-((4-chloro-phenyl)sulfonylamino carbonyl)phenylcarbonyl, and 4-((4-bromophenyl) sulfonylamino carbonyl) phenylcarbonyl; (ii) aromatic urethane protecting groups such as phenyloxycarbonyl, benzyloxycarbonyl and substituted benzyloxycarbonyls such as p-chlorobenzyloxycarbonyl, p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl, 1-(p-biphenyl)-1-methyl-ethoxycarbonyl, α,α -dimethyl-3,5-dimethoxybenzyloxycarbonyl, and benzhydryloxycarbonyl; (iii) aliphatic urethane protecting groups such as t-butyloxycarbonyl (Boc), diisopropylmethoxycarbonyl, isopropoxy-carbonyl, methoxycarbonyl, ethoxycarbonyl, isobutyloxycarbonyl and allyloxycarbonyl; (iv)

cycloalkyl urethane protecting groups such as cyclopentyloxycarbonyl, adamantyloxycarbonyl, and cyclohexyloxycarbonyl; (v) thiourethane protecting groups such as phenylthiocarbonyl; (vi) alkyl type protecting groups such as triphenylmethyl (trityl) and benzyl (Bn); and (vii) alkyl and aryl silyl protecting groups such as trimethylsilyl, tert-butyldimethylsilyl, tert-butyl-diphenylsilyl and triisopropylsilyl. t-butyloxycarbonyl (Boc) is one presently preferred α -amino protecting group. Representative methods using Boc for protecting are disclosed by Bodansky *et al.* in "The Practice of Peptide Synthesis," Springer-Verlag, Berlin (1984), p. 20). The selection and use of an appropriate coupling reagent is within the skill in the art. Particularly suitable coupling reagents where the amino acid to be added is Gln, Asn, or Arg include *N,N*-dicyclohexyl-carbodiimide and 1-hydroxybenzotriazole. The use of the latter reagents during synthesis of the instant compounds of FORMULA I, II and/or III prevents nitrile and lactam formation. Other suitable coupling agents include e.g., (i) carbodiimides such as 1-(3-dimethylamino-propyl)-3-ethyl-carbodiimide hydrochloride; (ii) ketenimines; (iii) isoxazolium salts such as N-ethyl-5-phenyl-isoxazolium-3-sulfonate; (iv) monocyclic nitrogen-containing heterocyclic amides of aromatic character containing 1-4 ring nitrogen atoms, e.g., imidazolides, pyrazolides and 1,2,4-triazolides, and particularly *N,N*-carbonyldiimidazole and *N,N*-carbonyl-di-1,2,4-triazole; (v) alkoxylated acetylene e.g., ethoxyacetylene; (vi) reagents which form a mixed anhydride with the carboxyl moiety of the amino acid, e.g., ethyl chloroformate and isobutyl chloroformate, or alternatively, the symmetrical anhydride of an amino acid to be coupled e.g., Boc-Ala-O-Ala-Boc; and (vii) nitrogen-containing heterocyclic compounds having a hydroxyl group on one ring nitrogen, e.g., N-hydroxyphthalimide, N-hydroxysuccinimide and 1-hydroxybenzotriazole. Activating reagents, and methods of their use in peptide coupling, are known in the art, e.g. as disclosed in Kapoor ((1970) *J. Pharm. Sci.* 59:1-27). For synthesis of the instant compounds according to FORMULAS I, II and/or III, presently preferred coupling agents are symmetrical anhydrides. For removal of a compound from a preferred benzhydrylamine resin, treatment with a solution of dimethyl sulfide, p-cresol, thiocresol, or anisole in anhydrous hydrogen fluoride is presently preferred.

The cleavage reaction is also preferably carried out at a temperature between about 0°C and about room temperature, over a period of about 5 minutes to about 5 hours.

Embodiments of the invention provide pharmaceutical compositions comprising one or more compounds according to FORMULA I, II and/or III, above. Preferably, each of the instant compounds present in the pharmaceutical composition is present in a therapeutically effective amount. The instant pharmaceutical compositions find a variety of uses in treatments for neurological disorders, e.g., schizophrenia, Alzheimer's disease, disorders of extrapyramidal motor function, such as Parkinson's disease, progressive supramuscular palsy, Huntington's disease, Gilles de la Tourette syndrome and tardive dyskinesia, obesity, severe pain, drug and tobacco withdrawal, respiration, mood and emotional disorders such as depression, anxiety and psychosis, motor control and function, focus and attention disorders, concentration disorders, memory loss, cognitive impairment, dementia (including AIDS dementia), neurodegenerative disorders, epilepsy; as well as in treatments of other diseases, e.g., cardiovascular dysfunction including hypertension and cardiac arrhythmias, convulsive disorders, eating disorders, including bulimia and anorexia, autonomic disorders including dysfunction of gastrointestinal motility and function such as inflammatory bowel disease, irritable bowel syndrome, diarrhea, constipation, gastric acid secretion and ulcers. The instant pharmaceutical compositions may also be useful as co-medication in surgical procedures and pheochromocytoma; or in co-treatments of heart insufficiency, hyperprolactinemia, bacterial infections, asthma or arthritis.

The instant pharmaceutical compositions are preferably formulated as solutions, suspensions, tablets, dispersible tablets, pills, capsules, powders, sustained release formulations or elixirs. As such the compositions are useful for oral administration, or in sterile solutions or suspensions, for parenteral administration. Transdermal delivery, e.g., patch preparations, and dry powder inhalers are also contemplated. Examples of methods that may be useful for formulating the instant pharmaceutical compositions are disclosed e.g., by Ansel, *Introduction to Pharmaceutical Dosage Forms, Fourth Edition* 1985, p. 126. The instant pharmaceutical compositions may be formulated as the sole pharmaceutically active

surfactants, e.g., TWEEN, dissolution in aqueous sodium bicarbonate, and preparation of soluble derivatives or prodrugs of the instant compounds prior to formulation.

Unit dosage forms suitable for administration to subjects in need thereof, e.g., as packaged individually, are known in the art. Each unit-dose commonly contains a predetermined quantity of the therapeutically active compound sufficient to produce a desired therapeutic effect; in association with the required pharmaceutical carrier, vehicle or diluent. Examples of unit-dose forms include ampoules and syringes and individually packaged tablets or capsules. Unit-dose forms may be administered in fractions or multiples thereof. Representative methods for preparing dosage forms are known in the art, e.g., see *Remington's Pharmaceutical Sciences*, Mack Publishing Company, Easton, Pa., 15th Edition, 1975.

Oral pharmaceutical dosage forms including solid, gel and liquid are contemplated formed as tablets, capsules, granules, bulk powders, chewable lozenges enteric-coated capsules, sugar-coated or film-coated capsules, hard or soft gelatin capsules, granules e.g., as non-effervescent or effervescent forms, pills, troches and the like. Liquid oral dosage forms include aqueous solutions, emulsions, suspensions, solutions and/or suspensions reconstituted from non-effervescent granules and effervescent preparations reconstituted from effervescent granules.

Parenteral injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Typically a therapeutically effective dosage is formulated to contain a concentration of at least about 0.1 % w/w up to about 90% w/w or more, preferably more than 1 % w/w of the active compound to the treated tissue(s).

Topical solutions, suspension, emulsions and the like may be formulated as creams, gels, ointments, emulsions, solutions, elixirs, lotions, suspensions, tinctures, pastes, foams, irrigations, sprays, nasal sprays, suppositories, bandages, dermal patches, transdermal delivery systems and the like. Contemplated formulations include inhalent aerosols in the form of an aerosol or nebulizer solutions, inhalent therapy solutions, microfine powders for insufflation, and the like. Also

contemplated are formulations for skin and mucous membranes, ophthalmic uses (e.g., eyes), intra-cisternal and intra-spinal application.

The pharmaceutical compositions containing compounds or pharmaceutically acceptable derivatives may be packaged as articles of manufacture containing e.g., a packaging material; the instant pharmaceutical composition; a product insert describing that the composition is effective in ameliorating one or more symptoms of disease, i.e., as set forth above; and, a label indicating that the composition containing the compound or derivative thereof is useful in the treatment or prevention of the disease, i.e., as set forth above.

Embodiments of the invention provide compounds according to FORMULA I, II and/or III that find uses in pharmaceutical compositions manufactured for the purpose of administering a treatment effective in altering apoptosis and altering processing of APP to A β , e.g., altering deposition of amyloid. The dose ranges, which can be established empirically, for use in the treatment of disease states will depend upon the etiology, nature, and severity of the disease state as well as such other factors as determined by the attending physician. The broad range for effective treatment is about 0.01 to 10 mg per kilogram (kg) of body weight per day. The preferred range is about 0.1 to 10 mg/kg of body weight per day. Preferred modes of administration include oral and parenteral modes of administration. It is believed that the instant treatments for patients with such amyloid disorders will effect beneficial changes in the disease process resulting in a stabilization of disease activity, slowed progression of disease, and/or an enhanced lifestyle with increased cognitive ability and decreased anxiety, as well as a possible delay (or obviation) of the need to institutionalize certain patients.

In other embodiments, pharmaceutical compositions are provided comprising the instant compounds of FORMULA I, II and/or III as formulated for use to alter apoptosis, e.g., by inhibiting the action of cysteinyl caspase proteases and/or cathepsin proteases. Ca²⁺ dependent proteases are presently thought to play significant roles in the pathology of a variety of disorders, including cerebral ischaemia, cataract, myocardial ischaemia, muscular dystrophy and platelet aggregation.


The following examples are included for illustrative purposes only and are not intended to limit the scope of the invention.

EXAMPLE 1

Solution Phase Synthesis:

(2*RS*)-2-*tert*-Butoxycarbonylamino-4-methyl-4-pentenoic Acid

A. Ethyl (2*RS*)-2-Diphenylmethylenamino-4-methyl-4-pentenoate



To a stirred solution of N-(diphenylmethylenamino)glycine ethyl ester (available from Aldrich Chemical Co., Milwaukee, WI) (50.0 g, 187 mmol) in dry THF (200 mL) at -78°C under argon was added a 1.0 M solution of lithium bis(trimethylsilyl)amide (Li HMDS; 210 mL, 0.2 mol) in THF over a period of 45 min ("1.A, Scheme A). After 0.5 hour at -78°C, 3-bromo-2-methylpropene (18.91 mL, 187 mmol) was added dropwise to the orange-yellow suspension. After the addition was complete, the reaction mixture was stirred for an additional 0.5 hour at 0°C and then allowed to warm to room temperature. After 1 hour the reaction mixture was concentrated under reduced pressure to give an orange-colored oil. The crude material was dissolved in ethyl acetate (EA) (600 mL), washed with brine (100 mL), dried over anhydrous MgSO₄ and concentrated under reduced pressure to give the


desired compound (60 g, 99%), which was used for the next step without further purification.

^1H NMR (CDCl_3 , 300 MHz), δ 7.1 - 7.7 (m, 10 H), 4.72 (dd, 2 H), 4.2 (m, 3H), 2.63 (m, 2H), 1.5 (s, 3H), 1.27 (t, 3 H).

B. Ethyl (2*RS*)-2-Amino-4-methyl-4-pentenoate hydrochloride

To a stirred solution of the alkylated Schiff's base from EXAMPLE 1.A above (60g, 0.187 mol) in diethyl ether (1.2 L) was added 1 N aqueous HCl (500mL). The resulting biphasic mixture was stirred for 24 hours at RT. The aqueous layer was separated from the ether layer, washed twice with EA (200 mL) and the aqueous solution was concentrated under reduced pressure to give the desired product "B", Scheme A (36 g, 99%) as viscous oil.

^1H NMR (CDCl_3 , 300 MHz) δ 5.1 (br s, 1 H), 4.9 (br s, 1 H), 4.32 (m, 3H), 2.5 - 2.9 (m, 2H), 1.83 (d, 3H), 1.36 (t, 3H).



C. Ethyl (2*RS*)-2-tert-Butoxycarbonylamino-4-methyl-4-pentenoate

To a stirred solution of ethyl (2*RS*)-2-amino-4-methyl-4-pentenoate "B", EXAMPLE 1.B, above, (6.0 g, 0.031 mol) in CH_2Cl_2 (75 mL) was added di-tert-butyl

dicarbonate (Boc)₂O; 6.79 g, 0.031 mol) followed by triethylamine (TEA, 8.7 mL, 0.062 mol) and the reaction mixture was left stirred at RT overnight. After overnight, the reaction mixture was diluted with water (100 mL), the organic layer was separated and washed with 0.5 N aqueous HCl (75 mL), water (75 mL, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give a yellow oil. The resultant crude material was purified via chromatography on silica (hexanes (H):EA, 3:1) to give ethyl (2*RS*)-2-*tert*-butoxycarbonylamino-4-methyl-4-pentenoate as a colorless oil (6.9 g, 87%).

¹H NMR (CDCl₃, 300 MHz) δ 4.9 (m, 1 H), 4.8 (br s, 1 H), 4.7 (br s, 1 H), 4.4 (m, 1 H), 4.2 (dq, 2 H), 2.3-2.6 (m, 2 H), 1.7 (s, 3 H), 1.42 (s, 9 H), 1.25 (t, 3 H).

D. (2*RS*)-2-*tert*-Butoxycarbonylamino-4-methyl-4-pentenoic Acid

To a stirred solution of ethyl (2*RS*)-2-*tert*-butoxycarbonylamino-4-methyl-4-pentenoate EXAMPLE 1.C, above (6.9 g, 26.9 mmol) in acetone (50 mL) was added LiOH (108 mL of 1 N aqueous solution, 108 mmol). The clear solution was stirred at room temperature for 0.5 hour. Acetone was evaporated under reduced pressure, the pH of the resulting aqueous solution was then adjusted to 3 by addition of 1 N HCl and the aqueous layer was extracted with EA (3 x 50 mL). The organic layers were combined, washed with brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give the desired acid as a pale yellow oil in quantitative yield.

¹H NMR (CDCl₃, 300 MHz) δ 4.95 (m, 1 H), 4.9 (br s, 1 H), 4.8 (br s, 1 H), 4.4 (m, 1 H), 2.3-2.7 (m, 2 H), 1.76 (s, 3 H), 1.55 (s, 9 H).

EXAMPLE 2

(3*RS*)-1-Phenylsulfonyl-3-amino-5-methyl-1E,5-hexadiene Hydrochloride

A. N,O-Dimethyl (2*RS*)-2-*tert*-Butoxycarbonylamino-4-methyl-4-pentenoylhydroxamate

To a solution of (2*RS*)-2-*tert*-butoxycarbonylamino-4-methyl-4-pentenoic acid (EXAMPLE 1C; Scheme B) in CH₂Cl₂ (50 mL) was added 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride (EDC) (5.67 g, 0.296 mol) and N,O-dimethylhydroxylamine hydrochloride (MeHNOMe.HCl; 2.89 g, 0.0296 mol).

The resulting suspension was cooled to 0°C, TEA (4.12 mL, 0.0296 mol) was added dropwise and the reaction mixture was allowed to warm to room temperature and left stirring overnight. After overnight, the reaction mixture was poured into cold water (75 mL) and extracted three times with CH₂Cl₂ (350 mL). The organic layers were combined, washed with 1 N HCl (75 mL), brine (75 mL), dried over Na₂SO₄ and concentrated under reduced pressure. The resultant crude product was purified by flash chromatography on silica gel using chloroform/methanol (99:1) as eluant to afford the desired compound (7.33 g, 53%).

¹H NMR (CDCl₃, 300 MHz) 5.2 (m, 1H), 4.83 (br s, 1H), 4.76 (br s, 1H), 3.8 (s, 3H), 3.2 (s, 3H), 2.1-2.5 (m, 2H), 1.77 (s, 3H), δ 1.4 (s, 9H).

B. (2*RS*)-2-tert-Butoxycarbonylamino-4-methyl-4-pentenal

Lithium aluminum hydride (LAH, 0.42 g, 11 mmol) was suspended in anhydrous diethyl ether (50 mL). A solution of the Weinreb amide in diethyl ether (10 mL) was added dropwise at 0°C. After 0.5 hour at this temperature, the reaction mixture was quenched by sequential addition of water (0.42 mL), 15% aqueous NaOH solution (0.42 mL) and water (1.26 mL). The precipitated inorganic salts were filtered through celite, and the filter pad washed with ether (50 mL). The filtrate was concentrated under reduced pressure to give the aldehyde as a clear viscous oil (3.35 g). The compound was used the same day for the subsequent steps without further purification.

C. Diethyl (Phenylsulfonyl)methylphosphonate

A stirred solution of methyl phenyl sulfone (20 g, 0.128 mol) in dry THF (200 mL) was cooled at -78°C in an acetone/dry ice bath. Lithium bis(trimethylsilyl)amide (Li HMDS; 134 mL, 1 M solution in THF, 0.134 mol) was added dropwise over 1 h. The resulting white suspension was stirred for an additional hour at -78°C and diethyl chlorophosphate (18.5 mL, 128 mmol) in THF (20 mL) was added. The reaction mixture was stirred at -78°C for 0.5 h, then allowed to slowly warm to room temperature and left stirring for 4h. The reaction mixture was next concentrated under reduced pressure, the residue was dissolved in EA (500 mL), washed with brine (100 mL), dried over anhydrous Na₂CO₃ and concentrated under reduced


pressure. The resultant crude material was purified by flash chromatography on silica gel (EA:H, 3:1 to 1:1) to afford the desired product as white solid (17.9 g, 48%).

¹H NMR (CDCl₃, 300 MHz) δ 7.5-8.1 (m, 5 H), 4.16 (q, 4 H), 3.8 (s, 1 H), 3.74 (s, 1 H), 1.29 (t, 6 H).

D. (3*RS*)-3-*tert*-Butoxycarbonylamino-5-methyl-1-phenylsulfonyl-1*E*,5-hexadiene

To a stirred solution of diethyl (phenylsulfonyl)methylphosphonate (from EXAMPLE 2.C) (3.54 g, 0.0121 mol) in dry THF (70 ml) at -78°C was added slowly a solution of lithium bis(trimethylsilyl)amide (LiHMDS; 13.2 ml, 1 M solution in THF, 13.2 mol) over 0.5 h. The reaction mixture was stirred for another 0.5 h at -78°C, and a solution of crude (2*RS*)-2-*tert*-butoxycarbonylamino-4-methyl-4-pentenal from EXAMPLE 2.B (2.35 g, 11.0 mol) in dry THF (20 ml) was added dropwise. After addition was complete, the reaction mixture was stirred for an additional 0.5 hour and then allowed to warm to room temperature and stirred for an additional 3 h. The stirred reaction mixture was concentrated under reduced pressure and the residue was dissolved in EA (100 mL). The EA solution was washed with brine (50 mL), dried over MgSO₄ and concentrated under reduced pressure to give an oil. The resultant crude material was purified via flash chromatography on silica gel (H:EA, 6:1) to provide the desired sulfone "D", Scheme B (1.82 g, 47% overall yield for 2.C and 2.D).

¹H NMR (CDCl₃, 300 MHz) δ 7.9-7.5 (m, 5 H), 6.9 (br d, 1 H), 6.44 (d, 1 H), 4.88 (br s, 1 H), 4.76 (br s, 1 H), 4.5 (m, 2 H), 2.2-2.3 (m, 2 H), 1.72 (s, 3 H), 1.37 (s, 9 H).



E. (3*RS*)-1-Phenylsulfonyl-3-amino-5-methyl-1*E*,5-hexadiene

Hydrochloride

To a stirred solution of the vinyl sulfone from EXAMPLE 2.D, above, (0.75 g, 2.13 mmol) in CH₂Cl₂ (10 ml) was added HCl (10.5 mL of a 4 M dioxane solution, 43.0 mol). The reaction mixture was stirred at room temperature for 0.5 h and then concentrated under reduced pressure to provide 0.61 g (99%) of the desired product as a white solid.

¹H NMR (CD₃OD, 300 MHz) δ 7.92-7.89 (m, 2 H), 7.72 - 7.61 (m, 3 H), 7.05- 6.83 (m, 2 H), 5.02 (br s, 1H), 4.85 (br s, 1 H), 4.21 (q, 1 H), 2.6-2.48 (m, 2 H), 1.71 (s, 3 H).

EXAMPLE 3

(3*RS*)-L-Leu-N-[3-(1-phenylsulfonyl)-5-methylhexa-1*E*,5-dieneamide

A. (3*RS*)-N-Boc-L-Leu-N-[3-(1-phenylsulfonyl)-5-methylhexa-

1*E*,5-dieneamide

To a stirred solution of N-Boc-Leu-OH (available from Aldrich Chemical Co., Milwaukee, WI) (1.66 g, 6.7 mmol) in anhydrous CH₂Cl₂ (30 mL) at RT were added 1-hydroxybenzotriazole (HOBt) (1.64 g, 12.1 mmol), EDC (1.28 g, 6.7 mmol), (3*RS*)-1-phenylsulfonyl-3-amino-5-methyl-1*E*,5-hexadiene hydrochloride from EXAMPLE 2.E (1.74 g, 6.06 mmol) and di-isopropylethylamine (DIEA) (0.86 g, 6.7 mmol). The reaction mixture was stirred at room temperature for 17 h and then concentrated under reduced pressure. The residue was resuspended in ethyl acetate (EA) (100 mL, washed successively with saturated bicarbonate solution (50 mL),

10% citric acid (50 mL), and brine (50 mL), dried over anhydrous MgSO_4 , and concentrated under reduced pressure. The resultant crude product, ("F, Scheme C), was purified by chromatography (EA:H, 1:2) to yield the desired product as colorless oil (2.19 g, 78%).

^1H NMR (CDCl_3 , 300 MHz) δ 7.87-7.80 (m, 2 H), 7.61-7.48 (m, 3 H), 6.9 (dt, 1 H), 6.5 (m, 1 H), 6.3 (m, 1 H), 6.15 (m, 1 H), 4.86 (br s, 1 H), 4.17 (br s, 1 H), 3.9 (m, 1 H), 2.2-2.4 (m, 2 H), 1.7 (s, 3H), 1.4 (br s, 9H), 0.9 (m, 9 H).

B . (3RS)-L-Leu-N-[3-(1-phenylsulfonyl)-5-methylhexa-1E,5-dieneamide]

This compound was prepared following the procedure of EXAMPLE 2.D and 2.E, above using (3RS)-3-tert-butyloxycarbonylamino-5-isobutyl-1-phenylsulfonyl-1E,5 hexadiene as a starting material.

^1H NMR (CD_3OD) δ 8.62-8.45 (m, 1 H), 7.77-7.70 (m, 2 H), 7.58-7.43 (m, 3 H), 6.80-6.71 (m, 1 H), 6.47-6.42 (m, 1 H), 4.67-4.61 (m, 2 H), 3.68-3.63 (m, 2 H), 2.25-2.00 (m, 2 H), 1.56-1.44 (s overlaps m, 6 H), 0.79 (d, 6 H).

EXAMPLE 4

(3RS)-N-Cbz-L-Val-L-Leu-N-[3-(1-phenylsulfonyl)-5-methyl-hexa-1E,5-dieneamide]

The title compound was prepared from N-Cbz-L-Val-OH (available from Bachem) and (3RS)-L-Leu-N-[3-(1-phenylsulfonyl)-5-methylenhexa-1E,5-dien]amide (EXAMPLE 3) using substantially the same procedure as described in EXAMPLE 3 to isolate the desired product as a colorless oil (10 mg).

^1H NMR (CDCl_3 , 300 MHz) δ 7.89-7.83 (m, 2 H), 7.60-7.5 (m, 3 H), 7.34-7.35 (m, 5 H), 6.92 (dd, 1 H), 6.56-6.51 (m, 1 H), 6.36 (m, 1 H), 5.32 (m, 1 H), 5.13-5.10 (m, 3 H), 4.85-4.75 (m, 3 H), 4.45 (m, 1 H), 3.93 (m, 1 H), 2.35 - 2.10 (m, 2 H), 1.73-1.47 (m, 7 H), 0.97-0.84 (m, 12 H).

EXAMPLE 5

**(3RS)-N-Cbz-L-Leu-L-Met-N-[3-O-phenylsulfonyl]-5-methylhexa-
-1E,5-dieneamide**


The title compound was prepared from N-Cbz-L-Leu-L-Met-OH synthesized by solid phase peptide synthesis (or purchased from Bachem) and using (3RS)-1-phenylsulfonyl-3-amino-5-methyl-1E,5-hexadiene hydrochloride (EXAMPLE 2.E) as a starting material in substantially the same procedure described in EXAMPLE 3 (Scheme C). The product was isolated as a colorless oil (10 mg).

^1H NMR (CDCl_3 , 300 MHz) δ 7.89-7.83 (m, 2 H), 7.60-7.47 (m, 3 H), 7.37-7.33 (m, 5 H), 6.97-6.90 (m, 2 H), 6.5 (m, 1 H), 5.28-5.23 (m, 4 H), 5.14-5.07 (m, 3 H), 4.52 (m, 1 H), 4.1 (m, 1 H), 2.5 - 1.64 (m overlaps s, 15 H), 0.92 (m, 6 H).

EXAMPLE 6

(2S)-2-tert-Butoxycarbonylamino-4-methyl-4-pentenoic Acid**A. (2S)-2-Amino-4-methyl-4-pentenoyl (S,S)-Pseudoephedrine Amide**

To a suspension of dry LiCl (4.2 g, 99 mmol) in dry THF (45 mL) was added diisopropylamine (DIA, 4.6 mL, 32.8 mmol). The flask was cooled to -78°C in a dry ice/acetone bath and a solution of n-BuLi (hexanes, 12.7 mL, 31.8 mmol) was added in drops. The solution was stirred for 10 minutes, then a solution of



(S,S)-pseudoephedrine glycinamide ("H", Scheme D; prepared according to Myers et al. *J. Am. Chem. Soc.* 117: 8488-8489, 1995) (3.6 g, 16.3 mmol) in THF (25 mL) was

added in drops over a period of 10 minutes. The reaction mixture was left stirring at -78°C for 20 minutes, changed to an ice bath and stirred at 0°C for 20 min. 3-Bromo-2-methylpropene (1.8 ml, 17.8 mmol) was added dropwise to the reaction mixture. After 0.5 h, 1.0 N HCl (110 mL) was added followed by EA (150 mL). The aqueous layer was separated and the EA solution was extracted with 1.0 N HCl (110 mL). The combined HCl solution was cooled in an ice bath and the pH adjusted to pH 14 using 50% NaOH. The basic solution was extracted twice with CH₂Cl₂ (80 ml each) and the combined CH₂Cl₂ solutions were dried over anhydrous K₂CO₃, filtered and concentrated to give a colorless oil that was purified by flash chromatography on silica gel (5% MeOH in CH₂Cl₂) to give a colorless oil (2.95 g, 66%).

¹H NMR (CDCl₃, 300 MHz) (mixture of rotomers) δ 7.38-7.27 (m, 5 H), 4.94-4.78 (m, 2 H), 4.65-4.46 (m, 2 H), 3.91-3.87 (dd, 0.2 H), 3.79-3.74 (dd, 0.8 H), (m, 1 H), 3.11-2.98 (br, 1 H), 2.97 (s, 0.6 H), 2.89 (s, 2.4 H), 2.22-1.74 (m overlaps s, 7 H), 1.12-1.09 (dd, 2.4 H), 1.01-1.08 (d, 0.6 H).

B. (2S)-2-tert-Butoxycarbonylamino-4-methyl-4-pentenoic Acid

To an aqueous solution of 1.0 N NaOH (15.2 ml, 15.2 mmol) was added a solution of (2S)-2-amino-4-methyl-4-pentenoyl (S,S)-pseudoephedrine amide (EXAMPLE 6.A, above) (2.03 g, 7.6 mmol) in water (15.2 mL). The reaction mixture was stirred under reflux for 1.5 h, and cooled to RT. The crystallized pseudoephedrine was filtered, and the filtrate was diluted with water (75 mL). The aqueous solution was extracted twice with CH₂Cl₂ (150 mL). The combined CH₂Cl₂ solutions were extracted with water (100 mL) and the aqueous solution was washed with CH₂Cl₂ (75 mL). The aqueous solutions were then combined and concentrated under reduced pressure, about 50 mL, solid NaHCO₃ (1.49 g) was added to this aqueous solution followed by p-dioxane (55 mL). The resulting solution was cooled to 0°C and di-tert-butyl dicarbonate (Boc₂O; 2.0 g, 9.17 mmol) was added. The reaction was left stirring at 0°C for 1 h, and the temperature was warmed to room temperature and stirred for an additional 0.5 h. The reaction mixture was diluted with water (150 mL) and extracted with EA (i.e., DIEA; 150 mL). The EA solution

was re-extracted with 2% NaHCO₃ (100 mL) solution. The combined bicarbonate solutions were then acidified to pH 3 using 10% citric acid solution. The acidic solution was extracted three times with EA (100 mL). The combined EA solutions were washed with water (2 times; 100 mL each), dried with MgSO₄, filtered and concentrated to give 1.46 g (84%) of product as colorless oil.

¹H NMR (CDCl₃, 300 MHz) δ 9.8 (br, 1 H), 5.3-4.80 (m, 3 H), 4.34-4.31 (m, 1 H), 2.62-2.56 (m, 2 H), 1.76 (s, 3 H), 1.43 (s, 9 H).

EXAMPLE 7

(3*S*)-N-Cbz-L-Leu-L-Leu-N-[3-(*l*-phenylsulfonyl)-5-methylhexa-1*E*,5-dieneamide

N-Cbz-L-Leu-L-Leu (Bachem), the (2*S*)-2-tert-butoxycarbonylamino-4-methyl-4-pentenoic acid product of EXAMPLE 6.B and diethyl-(phenylsulfonyl)-methyl-phosphonate (EXAMPLE 2.C) were combined using the procedure of EXAMPLES 2 and 3, above, to provide the title compound as a colorless oil.


¹H NMR (CDCl₃, 300 MHz) δ 7.86-7.83 (m, 2 H), 7.59-7.46 (m, 2 H), 7.41-7.29 (m, 5 H), 6.95-6.90 (dd, *J* = 3.9, 11 Hz, 1 H), 6.79 - 6.66 (br d, *J* = 3.9 Hz, 1 H), 6.59-6.56 (m, 2 H), 5.46-5.44 (br m, *J* = 6.9 Hz, 1 H), 5.14-4.99 (AB q, *J* = 12.3 Hz, 2 H), 4.84-4.72 (m, 3 H), 4.38-4.34 (m, 1 H), 4.13-4.09 (m, 1 H), 2.30-2.03 (m, 2 H), 1.67 (s, 3 H), 1.64-1.25 (m, 6 H), 0.89 - 0.81 (overlapping doublets, 12H).

EXAMPLE 8

(3*R*)-N-Cbz-L-Leu-L-Leu-N-[3-(*l*-phenylsulfonyl)-5-methylhexa-1*E*,5-dieneamide

The title compound was prepared according to the procedure set forth in EXAMPLE 7, using (2*R*)-2-tert-butoxycarbonylamino-4-methyl-4-pentenoic acid in place of (2*S*)-2-tert-butoxycarbonylamino-4-methyl-4-pentenoic acid, to afford the desired product as a colorless oil. (2*R*)-2-tert-butoxycarbonylamino-4-methyl-4-pentenoic acid was prepared according to EXAMPLE 6, above, using (*R,R*)-pseudoephedrine glycineamide.

EXAMPLE 9
(3RS)-4-Benzylpiperidinocarbonyl-L-Leu-L-Leu-N-[3-(1-phenylsulfonyl)-5-methylhexa-1 E,5-dienelamide



A. 4-Benzylpiperidinocarbonyl-L-Leu-OMe

To a stirred solution of triphosgene (1.02 g, 3.4 mmol) in anhydrous CH₂Cl₂ (50 mL) was added a solution of L-Leu-OMe (available from Aldrich Chemical Co., Milwaukee, WI) (1.8 g, 10.3 mmol) and TEA (1.43 mL, 10.3 mmol) in CH₂Cl₂ (25 mL) dropwise over a period of 15 min. (Scheme E). The reaction mixture was stirred at room temperature for 3 h and then a solution of 4-benzylpiperidine (1.81 mL, 10.3 mmol) and TEA (1.45 mL) in anhydrous CH₂Cl₂ (25 mL) was added in dropwise. The reaction mixture was then stirred at room temperature for 12 h, concentrated under reduced pressure. The residue was re-dissolved in EA (100 mL) and was washed successively with saturated bicarbonate (50 mL), brine (50 mL), 1.0 N HCl (50 mL) and saturated brine (50 mL), dried over anhydrous MgSO₄ filtered and concentrated to give a colorless oil that was purified by silica gel chromatography (2% MeOH:CH₂Cl₂) to yield the desired product as a colorless oil (2.6 g, 73%).

¹H NMR (CDCl₃, 300 MHz) δ 7.24-7.05 (m, 5H), 4.81-4.78 (d, J = 5.1 Hz, 1 H), 4.46-4.39 (m, 1 H), 3.91-3.83 (m, 2 H), 3.63 (s, 3 H), 2.71-2.59 (m, 2 H), 2.48-2.46 (d, 2 H), 1.65-1.40 (m, 6 H), 1.18 - 1.08 (m, 2 H), 0.88 - 0.86 (d, 6 H).

B. 4-Benzylpiperidinocarbonyl-L-Leu-OH

A solution of 4-benzylpiperidinocarbonyl-L-Leu-OMe (EXAMPLE 9.A., above) (2.6 g, 7.5 mmol) in MeOH (60 ml) was cooled to 0°C, LiOH·H₂O (1.26 g, 30 mmol) in water (20 mL) was added and the reaction mixture was stirred at 0°C for 2 h and at room temperature for 12 h (Scheme E). The reaction mixture was then concentrated under reduce pressure to 30 ml, diluted with EA (100 ml) and the EA solution was acidified with 1.0 N HCl to pH 1. The organic layer was separated and the aqueous layer was extracted with EA (50mL). The combined EA solutions were washed with water (2 time; 50 mL), dried over anhydrous MgSO₄ filtered and concentrated under reduced pressure to give the desired product as a colorless oil (2.5 g, 99%).

¹H NMR (CDCl₃, 300 MHz) 7.32-7.11 (m, 5 H), 5.26-5.23 (d, 1 H), 4.53-4.48 (m, 1 H), 3.94-3.89 (m, 2 H), 2.74-2.66 (br t, 2 H), 2.54-2.52 (d, 2 H), 1.86-1.52 (m, 6 H), 1.32-1.22 (m, 2 H), 0.93 (d, 6 H).

C. 4-Benzylpiperidinocarbonyl-L-Leu-L-Leu-OMe

The title compound was prepared using the procedure set forth in EXAMPLE 9.A, above, using Leu-Leu-OMe as a starting material instead of Leu-OMe. The product was isolated as a colorless oil (332 mg, 69%).

¹H NMR (CDCl₃, 300 MHz) δ 7.48-7.1 (m, 5 H), 5.28-5.21 (br d, 1 H), 4.61-4.40 (m, 2 H), 3.94-3.82 (m, 2 H), 2.71 (t, 2 H), 2.76 (d, 2 H), 1.90-1.69 (m, 9 H), 1.48-1.34 (m, 2 H), 1.17 - 1.12 (m, 12 H).

D. 4-Benzylpiperidinocarbonyl-L-Leu-L-Leu-OH

The title compound was prepared as a colorless oil from the product of 9.C using the procedure of EXAMPLE 9.B.

¹H NMR (CDCl₃, 300 MHz) δ 7.34-7.11 (m, 5 H), 5.24 (d, 1 H), 4.61-4.48 (m, 3 H), 3.94-3.90 (br m, 2 H), 2.70 (br t, 2 H), 2.53 (d, 2 H), 1.71-1.59 (m, 9 H), 1.24-1.10 (m, 2 H), 0.96-0.88 (m, 12 H).

E. (3RS)-4-Benzylpiperidinocarbonyl-L-Leu-L-Leu-N-[3-(1-phenylsulfonyl)-5-methylhexa-1E,5-dieneamide]

To a stirred solution of 4-benzylpiperidinocarbonyl-L-Leu-L-Leu-OH (138 mg, 0.31 mmol) in 3 mL of anhydrous CH_2Cl_2 at room temperature was added HOBt (42 mg, 0.31 mmol), EDC (65 mg, 0.34 mmol), DIEA (107 mL, 0.62 mmol) and 3(RS)-1-phenylsulfonyl-3-amino-5-methyl-1E,5-hexadiene hydrochloride from EXAMPLE 2.E (Scheme E). The reaction mixture was stirred at room temperature for 12 h, diluted with 50 mL of EA, washed with 50 mL of saturated bicarbonate solution, 10% citric acid and water, dried with MgSO_4 , filtered and concentrated to give crude product that was purified by silica gel chromatography (3% MeOH in CH_2Cl_2) to yield 170 mg (81 %) of product as white foamy solid. ^1H NMR (CDCl_3 , 300 MHz) δ 7.89 - 7.79 (m, 2 H), 7.61-7.45 (m, 3 H), 7.31- 7.11 (m, 5 H), 6.95-6.87 (m, 1 H), 6.76-6.72 (m, 1 H), 6.61-6.48 (m, 1 H), 4.96 (d, 1 H), 4.87-4.65 (m, 3 H), 4.35-4.25 (m, 1 H), 4.16-3.80 (m, 3 H), 2.87-2.85 (m, 2 H), 2.65 (d, 302 H), 2.36-2.20 (m, 3 H), 1.79-1.40 (m, 12 H), 1.21-1.07 (m, 2 H), 0.90- 0.79 (m, 12 H).


EXAMPLE 10

(3RS)-Tetrahydroisoquinolinylcarbonyl-L-Leu-L-Leu-N-[3-0-phenylsulfonyl)-5-methylhexa-1E,5-dieneamide]

The title compound was obtained as a glassy solid following the procedure described in EXAMPLE 9.A and 9.B, above, but using 1,2,3,4-tetrahydroisoquinoline as a reagent instead of 4-benzylpiperidine (Scheme F).

^1H NMR (CDCl_3 , 300 MHz) δ 7.85-7.83 (d, 2 H), 7.57-7.44 (m, 3 H), 7.27-

Scheme F



7.13 (m, 5 H), 6.96-6.89 (dd, 1 H), 6.85-6.83 (d, 1 H), 6.60-6.54 (dd, 1 H), 5.03-5.01 (d, 1 H), 4.84-4.73 (m, 3H), 4.63-4.46 (q, 2 H), 4.36-4.28 (m, 1 H), 4.24-4.18 (m, 1 H), 3.69-3.47 (m, 2 H), 2.89-2.84 (m, 2 H), 1.74-1.72 (d, J = 5.1 Hz, 2 H), 1.70 (s, 3 H), 1.74-1.46 (m, 6 H), 0.97-0.89 (overlapping doublets, 12 H).

EXAMPLE 11

(3RS)-Tetrahydroisoquinolinylicarbonyl-L-Val-L-Met-N-[3-(1-phenylsulfonyl)-5-methylhexa-1E,5-dieneamide]

The title compound was isolated as a colorless oil (100 mg) following the procedure outlined in EXAMPLE 10 (Scheme F), but using L-Val-OMe and L-Met-OMe as reagents (both available from Aldrich Chemical Co., Milwaukee, WI) instead of L-Leu-OMe.

¹H NMR (CDCl₃, 300 MHz) δ 7.85-7.80 (m, 2 H), 7.60-7.40 (m, 3 H), 7.19-7.12 (m, 5 H), 6.75 (dd, 1 H), 6.43-6.32 (m, 1 H), 5.14-4.44 (m, 8 H), 3.97-3.92 (m, 1 H), 3.70-3.50 (m, 2H), 2.87-2.84 (m, 2 H), 2.5-1.95 (m, overlaps s, 10 H), 1.7 (s, 3 H), 0.98-0.83 (m, 6 H).

EXAMPLE 12

(3RS)-Tetrahydroisoquinolinylicarbonyl-L-Val-L-Leu-N-[3-(1-phenylsulfonyl)-5-methylhexa-1E,5-dienelamide]

The title compound was isolated as a colorless oil (50 mg) following the procedure outlined in EXAMPLE 10 (Scheme F), above, but using L-Leu-OMe as a reagent in place of L-Met-OMe.

¹H NMR (CDCl₃, 300 MHz) δ 7.82-7.7 (m, 2 H), 7.55-7.45 (m, 3 H), 7.22-7.12 (m, 4 H), 7.09-7.01 (m, 1 H), 6.95-6.93 (m, 1 H), 6.62-6.12 (m, 2 H), 5.14-4.49 (m, 6 H), 4.36-4.20 (m, 1 H), 4.0-3.92 (m, 1 H), 3.71-3.53 (m, 2 H), 2.97-2.85 (m, 2 H), 2.49-2.02 (m, 3 H), 1.64-1.24 (m overlaps s, 6 H), 1.0-0.81 (m, 12 H).

EXAMPLE 13

(3S)-4-N-Benzylpiperazinylcarbonyl-L-Leu-L-Leu-N-[3-(1-phenylsulfonyl) hexamide]

Following the procedure set forth in EXAMPLE 9 (Scheme D), above, substituting 1-benzylpiperazine for 4-benzylpiperidine and substituting (3S)-1-phenylsulfonyl-3-amino-5-methyl-1E,5-hexadiene hydrochloride (e.g., prepared in a manner similar to that set forth in EXAMPLE 7) for (3RS)-1-phenylsulfonyl-3-

amino-5-methyl-1E,5-hexadiene hydrochloride. The title compound was isolated as a colorless oil.

¹H NMR (CD₃OD, 300 MHz) δ 7.79-7.77 (m, 2 H), 7.60-7.46 (m, 3 H), 7.24-7.16 (m, 5 H), 6.84-6.77 (dd, J = 5.1, 15 Hz, 1 H), 6.55-6.49 (dd, J = 1.5, 15 Hz, 1 H), 4.46-4.43 (m, 1 H), 4.21-4.17 (m, 1 H), 4.05-4.00 (m, 1 H), 3.40-3.20 (m, 6 H), 2.35-2.32 (m, 4 H), 1.62-1.39 (m, 8 H), 1.23 - 1.19 (m, 4 H), 0.88-0.77 (m, 15 H).

EXAMPLE 14

Illustrative *In Vitro* Assays

Summary Overview: Illustrative methods are described above by which the instant compounds may be synthesized. Testing the intracellular effects of a compound may be accomplished according to cell-based methods disclosed herein, such as those illustrated below using cultured human glioblastoma (HGB) cells stably-transfected with cDNA constructs encoding APP, e.g., the 695 amino acid β -amyloid precursor protein (APP₆₉₅), as well as other possible truncated forms of APP.

Cell-Based Screening Assays: An *in vitro* system may be used to assay the instant compounds and confirm those that modulate APP:A β processing. For instance, the human glioblastoma (HGB) cell line normally secretes only soluble derivatives of APP₇₅₁, but APP:A β processing may be altered to resemble processing in Alzheimer's disease by stable transfecting the HGB cells with a cDNA construct encoding either: (i) a wild-type 695 amino acid APP (APP_{695wt}), or (ii) a 695 amino acid APP containing a "Swedish" double mutation with Asn for Lys and Leu for Met substitutions at codons 670 and 671, respectively, (APP_{670/671}), (codons numbered according to the 770 amino acid APP isoform), or (iii) a 695 amino acid APP containing the 670/671 double mutation and a Phe for Val substitution at codon 717 (APP_{670/671/717}). Altered APP:A β processing in the HGB cells may e.g. be determined by collecting conditioned medium from the stably transfected cells and immunoblotting with monoclonal antibody specific for an N-terminal sequence in APP (e.g., mAb22C1), and/or a monoclonal antibody specific for the amino terminus of A β (e.g. mAb6E10). One or more of the instant compounds are added to the transfected/alterd HGB cells and APP:A β processing determined by immunoblotting

with comparison being made between cell cultures, so treated, and untreated control cultures, i.e., using medium collected from the cell cultures and/or cell extracts.

M-17 neuroblastoma cells transfected with a mutant APP construct (APP_{695NL+I}) secrete high levels of A β and a significant fraction is A β ₁₋₄₂ (39). HGB cells stably-transfected with a construct encoding APP_{670/671/717} or APP_{695NL+F} will secrete detectable levels of an ~4 kD peptide, i.e., A β ₁₋₄₀ detectable by immunoprecipitations, SDS-PAGE and/or western blotting. The ability of the instant compounds to modulate APP:A β processing in these cells is detected by evaluating A β levels in the medium collected from cell cultures or in cell culture extracts.

Quantitative assays for A β can be used to measure the levels of APP processing to A β , both *in vitro* e.g., in one or more of the cell-based assays, above, and also in extracts of animal tissues or in primary cultures of mixed brain cells (termed mixed brain cultures). For example, immunoblotting of SDS-PAGE gels with polyclonal rabbit anti-APP₆₄₅₋₆₉₇ (i.e., R369) may be used to reveal the different molecular weight species of APP antigens in tissues. Immunoblotting with monoclonal antibody 26D6, specific for amino acid residues 1-12 of A β ₁₋₄₀, may be used to confirm the presence of A β in the tissue extracts. Materials and Methods are set forth following the EXAMPLES section, below.

Using such cell-based assays the following results are recorded *in vitro*:

1. Examples of the instant compounds according to FORMULAS I, II and/or III, above, decrease processing of APP to A β *in vitro*, i.e., as judged by decreased levels of A β in treated HGB cell extracts by ELISA, immunoprecipitation and SDS-PAGE and/or western blotting (TABLE I, next page);
2. The instant compound inhibit A β formation *in vitro* in HGB cells without altering the amount of non-APP proteins (e.g., fibronectin, fodrin etc.) synthesized, i.e., as judged by ³⁵S-biosynthetically radiolabeling proteins synthesized by cells on SDS-PAGE and autoradiography; and
3. The instant compounds inhibited certain cell free cysteinyl proteases *in vitro*, including Cathepsin S, Calpain I, papain, Cathepsin L, but not Chymotrypsin, Cathepsin H, Cathepsin B or Cathepsin C (TABLE II, below).

TABLE I
Inhibition of APP Processing to A β *

CMPD/ Formula ^a	R ₇	Q	n	R ₆	R ₅	R ₄	R ₃	R ₂	R ₁	R ₈	X	Y	Z	Aβ Inh.
1/I	benzyl	-COC(O)-	1	H	(CH ₃) ₂ C-	H	indolyl	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	4+
2/I	piperidinyl- benzyl-	-C(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₃) ₂ C	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	5+
3/I	piperidinyl- benzyl-	-C(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₃) ₂ C	H	butyl	H	-C=C-	-S(O) ₂ -	benzyl	1+
4/I	benzyl-	-COC(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₃) ₂ C	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	6+
5/I	benzyl-	-COC(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₃) ₂ C	H	butyl	H	-C=C-	-S(O) ₂ -	benzyl	2+
6/I	(CH ₃) ₃ C-	-OC(O)-	1	C H ₃	(CH ₃) ₂ C-	H	(CH ₃) ₂ C	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	2+
7/I	(CH ₃) ₃ C-	-OC(O)-	1	C H ₃	benzyl- CH ₂ -	H	(CH ₃) ₂ C	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	2+
8/I	benzyl	-COC(O)-	1	H	CH ₃ -	H	(CH ₃) ₂ C	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	2+
9/I	piperidinyl-	-C(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₃) ₂ C	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	7+
10/I	benzyl-	-COC(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₂) ₂ SCH ₃	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	8+
11/I	benzyl-	-COC(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₃) ₂ C-	H	butyl	H	-C=C-	-S(O) ₂ -	-CH ₃	9+
12/I	benzyl-	-COC(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₃) ₂ C-	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	11+
13/I	benzyl-	-COC(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₃) ₂ C-	H	iso- butenyl	H	-C=C-	-S(O) ₂ - benzyl	ethynyl	3+
14/I	benzyl-	-COC(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₃) ₂ C-	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	3+
15/I	benzyl- CH ₂ - piperazinyl	-C(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₂) ₂ SCH ₃	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	3+
16/I	benzyl	-COC(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₃) ₂ C- CH ₃	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	3+
17/I	benzyl- piperidinyl	-C(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₃) ₂ C-	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	3+
18/I	benzyl- piperidinyl	-C(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₂) ₂ SCH ₃	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	3+
19/I	benzyl	-COC(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₃) ₂ C-	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	12+
20/I	benzyl- CH ₂ - piperidinyl	-C(O)-	1	H	(CH ₃) ₂ C-	H	(CH ₃) ₂ C-	H	iso- butenyl	H	-C=C-	-S(O) ₂ -	benzyl	13+

a. Form.= FORMULA I, II or III, supra; *Inhibition of APP processing to A β determined according to the ELISA methods, *Materials and Methods*.

TABLE II
Inhibition of Protease Activity *In Vitro**

CMPD ^a	CatS	CatL	Calpain I	Papain	CatB	CatC	CatH	Chymo.
1	ND	3+	-	4+	-	-	-	-
2	ND	-	2+	-	-	-	-	-
4	ND	4+	-	4+	-	-	-	-
5	ND	-	2+	1+	-	-	-	-
8	ND	4+	1+	1+	-	-	ND	-
9	4+	-	3+	-	-	-	-	-
10	ND	3+	-	4+	-	-	-	-
12	-	-	-	-	-	-	-	-
14	4+	4+	2+	-	-	-	ND	-
20	4+	ND	ND	ND	ND	ND	ND	ND

* Fluorescent and/or colorimetric substrate assays using peptidyl substrates; ND, not determined; Cat, cathepsin, Chymo., chymotrypsin;

a) CMPD, compounds as recited in TABLE I, above.

EXAMPLE 15

Illustrative Results in Transgenic Animal Models

Several useful transgenic animal test models of Alzheimer's disease have been described, e.g. Tg2576 and APP-YAC transgenic mice. These animal models are useful for determining the modulatory effects of the instant compounds on APP processing to A β . For example, the levels of inhibition of A β production *in vivo* may be determined by measuring the levels of A β in brain tissue, CSF fluid or plasma using an ELISA, immunoprecipitation and SDS-PAGE and/or western immunoblot analyses.

Similarly, using APP-YAC homozygous transgenic mice, the instant compounds may be tested to determine whether they inhibit A β production. Using such assays the following results may be obtained: namely,

1. The instant compounds preferably inhibit A β production as judged by decreased plasma or brain tissue levels of A β after administration of a test dose of one of the instant compounds; and,
2. The instant compounds preferably inhibit A β production in brain tissues after administration, i.e., as judged by decreased levels of biosynthetically radiolabeled A β in immunoprecipitates from brain tissues.

Materials and Methods

Cell Lines: The HGB₆₉₅ cell line may be prepared by stably-transfecting human glioblastoma (U138 MG) cells with an APP₆₉₅ expression plasmid. A human neuroblastoma

cell line (IMR-32) is potentially useful as the source for mRNA encoding the sequence of APP₆₉₅.

APP₆₉₅ cDNA and plasmids: cDNA encoding the APP₆₉₅ may be derived from mRNA purified from human neuroblastoma cell line (IMR-32), e.g., by RT-PCR amplification followed by cloning into a retroviral construct. The DNA fragment containing the APP₆₉₅ sequence may be removed from a selected cloned retroviral construct, e.g., by partial digestion with endonuclease, and ligated into a mammalian expression vector (e.g. obtained from Invitrogen, Inc., San Diego, CA.)

A β Cells: Transfection of HGB (or other) cells with the APP₆₉₅ cDNA (above) may be accomplished using the Lipofectin kit followed by selection of transformant clones in G418-containing medium.

APP Cells: An expression plasmid construct containing APP₆₉₅ cDNA may be mutated using site-directed mutagenesis in conjunction with PCR to generate an APP₆₉₅ plasmid containing missense mutations. This approach has been used by others to yield APP₆₉₅ coding region mutations, e.g., at codons 670,671 and 717 (codon numbering based upon the APP₇₇₀ isoform numbering system), where it has proved useful to generate Asn for Lys, Leu for Met and Phe for Val substitutions, respectively. Resulting constructs of this type, e.g. APP_{670,671,717} may be useful for selecting stably-transfect cell lines. Stable cell populations may be subcloned; characterized for expression of exogenous APP₆₉₅ by immunoblotting with APP specific antibodies; and then selected for use in one or more cell-based assays.

Antibodies: The mouse monoclonal antibody, 26D6, anti-human A β ₁₋₁₂ was prepared by TSD Bioservices. 26D6 IgG, i.e., specific for the N-terminus of A β , i.e., A β ₁₋₁₂, was purified from ascites by affinity purification on Protein A-Sepharose (Pharmacia) chromatography followed by affinity chromatography on an affinity matrix, (i.e., CnBr-activated Sepharose), containing bound A β ₁₋₁₂, i.e., at an input concentration of 1 mg/ml resin. Affinity purified 26D6 was coupled to horse-radish peroxidase (HRP) by Taconic Bioservices (Newark, DE). Rabbit antiserum, i.e., R6257, was raised to APP₆₅₇₋₆₇₆.

ELISA for A β : An A β sandwich ELISA was developed using polyclonal rabbit anti-A β ₁₋₄₀ as a selective capture antibody and HRP-conjugated monoclonal 26D6 (i.e., mouse monoclonal IgG specific for the N-terminus of A β , i.e., A β ₁₋₁₂,) for the detect reagent. To measure intracellular levels of A β and APP, lysates of HGB_{695WT} cells may be

prepared, separated by 16% Novex gradient gels, and immunoblotted with R369, (i.e., rabbit CTF-specific antisera), or the 26D6 monoclonal antibody.

For example, APP α - and β -secretase carboxy-terminal products (α -CTF and β -CTF) may be immunoprecipitated with anti-CTF antisera (and Protein A or Protein G Sepharose), immunoprecipitates can be separated by SDS-PAGE, then A β detected by western blotting with quantification being achieved using laser scanning densitometry.

For detection of A β in plasma and CSF, an ELISA assay may be used, e.g., using mouse monoclonal IgG anti-A β ₁₋₄₀ as a capture antibody and biotinylated mouse monoclonal antibody 4G8 (i.e., anti-A β ₁₇₋₂₄) as a detect reagent. After washing to remove unbound detect reagent, avidin-HRP is added to generate a colorimetric signal in the assay.

Treatment Studies and Immunochemical Assays *In Vitro*: Identical confluent cultures of stably-transfected APP_{695wt} or APP_{670/671/717} cells (approximately 10⁷ cells/plate) can be established in 100 mm² culture dishes and washed with 5 ml of Ultraculture medium. Compounds of interest can be dissolved, e.g., in DMSO, and then diluted using medium. As a control, plates can be treated with medium and diluted DMSO vehicle. After about 16 h incubation at 37°C with test compound or vehicle, conditioned medium can be collected and clarified, e.g., by low speed centrifugation (1000 x g). Clarified conditioned medium (5 ml) samples can then be pre-cleared e.g., using non-immune rabbit sera and Protein A-Sepharose. Pre-cleared supernatants are immunoprecipitated using e.g. a monospecific rabbit anti-A β ₁₋₂₈ antibody (rpAb1-28) and Protein A-Sepharose. The samples can then be centrifuged (1000 x g) and the supernatants discarded. A β immunoprecipitates can be solubilized and separated on SDS-PAGE where the separated proteins on the gels can be transferred to nitrocellulose (Hybond) and western immunoblotted using a monoclonal antibody, e.g., 6E10 (Senetek, Maryland Heights, MO., ascites fluid diluted 1:3000). Proteins binding A β -antibodies can be identified using biotin-conjugated goat-anti-mouse IgG (Sigma, St. Louis, MO., 1:1000) and streptavidin-horseradish peroxidase conjugate (1:1000, Amersham, Arlington Heights, IL). The peroxidase activity may be visualized e.g. using chemiluminescence (Amersham ECL kit).

Compound toxicity may be assessed e.g. using a LIVE/DEAD EukoLight™ Viability/Cytotoxicity assay (Molecular Probes, Inc., Eugene, OR.) according to manufacturer recommended protocols.

Drug Treatment Studies *In Vivo*: For *in vivo* studies, APP-transgenic mice may be dosed, (e.g., subcutaneously, or, orally), with vehicle, or one or more of the instant

compounds (e.g., at 10, 50, 100 or 200 μ moles/kg; with about 8-9 animals per group per timepoint). Blood and brain tissues may be collected at different times after injection. Samples are subject to ELISA assay, i.e., for analysis of levels of $A\beta_{1-40}$ in plasma or in CHAPS extracted brain tissues. Immunoprecipitation (with 26D6 monoclonal antibody and anti-mouse IgG or Protein A), western blotting and gel scanning densitometry may be used to confirm $A\beta$ fragment levels in tissues and plasma.

Citations

1. Khachaturian, Z. S. (1985) *Arch. Neurol.* 42: 1097-1105.
2. Glenner, G. G., and Wong, C. W. (1984) *Biochem. Biophys. Res. Commun.* 120: 885-890.
3. Masters, C. L., Multhaup, G., Sims, G., Pottgiesser, J., Martins, R. N., and Beyreuther, K. (1985) *EMBO J.* 4: 2757-2763.
4. Kang, J., Lemaire, H.-G., Unterbeck, A., Salbaum, J. M., Masters, C. L., Grzeschik, K.-H., Multhaup, G., Beyreuther, K., and Muller-Hill, B. (1987) in *Nature* 325: 733-736.
5. Weidemann, A., König, G., Bunke, D., Fischer, P., Salbaum, J. M., Masters, C. L., and Beyreuther, K. (1989) *Cell* 57: 115-126.
6. Mattson, M.P. 1997. Cellular actions of beta-amyloid precursor protein and its soluble and fibrillogenic derivatives. *Physiol. Rev.* 77(4): 1081-1132.
7. Jarrett, J. T., Berger, E. P., and Lansbury, P. T. (1993) *Biochemistry* 32: 4693-4697.
8. Yankner, B. A., Duffy, L. K., and Kirschner, D. A. (1990) *Science* 245: 417-420.
9. Loo, D. T., Copani, A. G., Pike, C. J., Whittemore, E. R., Walencewicz, A. J., and Cotman, C. W. (1993) *Proc. Natl. Acad. Sci., U.S.A.* 90: 7951-7955.
10. Hensley, K., Carney, J. M., Mattson, M. P., Aksenova, M., Harris, M., Wu, J. F., Floyd, R. A., and Butterfield, D. A. (1994) *Proc. Natl. Acad. Sci., U.S.A.* 91: 3270-3274.
11. Behl, C., Davis, J. B., Lesley, R., and Shubert, D. (1994) in *Cell* 77: 817-827.
12. Rojers, J., Cooper, N. R., Webster, S., Schultz, J., McGeer, P. L., Styren, S. D., Civin, W. H., Brachova, L., Bradt, B., Ward, P., and Lieberburg, I. (1992) *Proc. Natl. Acad. Sci., U.S.A.* 89: 10016-10020.
13. Strittmatter, W. J., Weisgraber, K. H., Huang, D. Y., Dong, L.-M., Salvesen, G. S., Pericak-Vance, M., Schmechel, D., Saunders, A. M., Goldgaber, D., and Roses, A. D. (1993) *Proc. Natl. Acad. Sci., U.S.A.* 90: 8098-8102.
14. Schmechel, D. E., Saunders, A. M., Strittmatter, W. J., Crain, B. J., Hulette, C. M., Joo, S. H., Pericak-Vance, M. A., Goldgaber, D., and Roses, A. D. (1993) *Proc. Natl. Acad. Sci., U.S.A.* 90: 9649-9653.
15. Rebeck, J. W., Reiter, J. S., Strickland, D. K., and Hyman, B. T. (1993) *Neuron* 11: 575-580.
16. Sisodia, S. S., Koo, E. H., Beyreuther, K., Unterbeck, A., and Price, D. L. (1990) *Science* 248: 492-495.
17. Esch, F. S., Keim, P. S., Beattie, E. C., Blacher, R. W., Culwell, A. R., Oltersdorf, T., McClure, D., and Ward, P. J. (1990) *Science* 248: 1122-1124.
18. Estus, S., Golde, T. E., Kunishita, T., Blades, D., Lowery, D., Eisen, M., Usiak, M., Qu, X., Tabira, T., Greenberg, B., and Younkin, S. G. (1992) *Science* 255: 728.
19. Golde, T. E., Estus, S., Younkin, L., Selkoe, D. J., and Younkin, S. G. (1992) in *Science* Vol. 255: 728-730.
20. Munoz, B., Kounnas, M. Z., Srinivasan, K., Vernier, J.-M., Wagner, S. L., and McDonald, I. A. (1997) *Alzheimer's Disease ID Drug Alert* 1: 1-8.
21. Cole, G. M., Huynh, T. V., and Saitoh, T. (1989) *Neurochem. Res.* 14: 933-939.
22. Norstedt, C., Caporaso, G. L., Thyberg, J., Gandy, S. E., and Greengard, P. (1993) *J. Biol. Chem.* 268: 608-612.
23. Felsenstein, K. M., Hunihan, L. W., and Roberts, S. B. (1994) in *Nature. Genet.* 6: 251-256.
24. Barrow, C. J., and Zagorski, M. G. (1991) in *Science* 253: 179-182.
25. Burdick, D., Soreghan, B., Kwon, M., Kosmoski, J., Knauer, M., Henschen, A., Yates, J., Cotman, C., and Glabe, C. (1992) in *Journal of Biological Chemistry* Vol. 267: 546-554.
26. Haass, C., Hung, A. Y., Schlossmacher, M. G., Teplow, D., and Selkoe, D. J. (1993) *J. Biol. Chem.* 268: 3021-3024.
27. Busciglio, J., Gabuzda, D. H., Matsudaira, P., and Yankner, B. A. (1993) in *Proc. Natl. Academy of Science, U.S.A.* Vol. 90: 2092-2096.
28. Siman, R., Mistretta, S., Durkin, J. T., Savage, M. J., Loh, T., Trusko, S., and Scott, R. W. (1993) *J. Biol. Chem.* 268: 16602-16609.
29. Klafki, H.-W., Paganetti, P. A., Sommer, B., and Staufenbiel, M. (1995) *Neuroscience Letter* 201: 29-32.
30. Klafki, H. W., Abramowski, D., Swoboda, R., Paganetti, P. A., and Staufenbiel, M. (1996) *J. Biol. Chem.* 271: 28655-28659.




SD714

31. Higaki, J., Quon, D., Zhong, Z., and Cordell, B. (1995) in *Neuron* 14: 651-659.
32. Citron, M., Diehl, T.S., Capell, A., Haass, C., Teplow, D., and Selkoe, D.J. (1996) *Neuron* 17: 171-179.
33. Goate, A., Chartier-Harlin, M.-C., Mullan, M., Brown, J., Crawford, F., Fidani, L., Giuffra, L., Haynes, A., Irving, N., James, L., Mant, R., Newton, P., Rooke, K., Roques, P., Talbot, C., Pericak-Vance, M., Roses, A., Williamson, R., Rossor, M., Owen, M., and Hardy, J. (1991) *Nature* 349: 704-706.
34. Chartier-Harlin, M. C., Crawford, F., Houlden, H., Warren, A., Hughes, D., Fidani, L., Goate, A., Rossor, M., Roques, P., Hardy, J., and Mullan, M. (1991) *Nature* 353: 844-846.
35. Murrell, J., Farlow, M., Ghetti, B., and Benson, M. (1991) in *Science* 254: 97-99.
36. Mullan, M., Crawford, F., Axelman, K., Houlden, H., Lilius, L., Winblad, B., and Lannfelt, L. (1992) *Nature Genet.* 1: 345-347.
37. Citron, M., Oltersdorf, T., Haass, C., McConlogue, L., Hung, A. Y., Seubert, P., Vigo-Pelfrey, C., Lieberburg, I., and Selkoe, D. J. (1992) *Nature* 360: 672-674.
38. Cai, X.-D., Golde, T. E., and Younkin, S. G. (1993) in *Science* 259: 514-516.
39. Suzuki, N., Cheung, T. T., Cai, X.-D., Odaka, A., Otvos, L., Eckman, C., Golde, T. E., and Younkin, S. G. (1994) *Science* 264: 1336-1340.
40. Haass, C., Hung, A. Y., Schlossmacher, M. G., Teplow, D., and Selkoe, D. J. (1993) *J. Biol. Chem.* 268: 3021-3024.
41. Knops, J., Suomensari, S., Lee, M., McConlogue, L., Seubert, P., and Sinha, S. (1995) *J. Biol. Chem.* 270: 2419-2422.
42. Pike, C. J., Burdick, D., Walencewicz, A. J., Glabe, C. G., and Cotman, C. W. (1993) *J. Neuroscience* 13: 1676-1687.
43. Busciglio, J., Lorenzo, A., Yeh, J., and Yankner, B. A. (1995) in *Neuron* Vol. 14, pp. 879-888.
44. Meda, L., Cassatella, M. A., Szendrei, G. I., Otvos, L., Baron, P., Villalba, M., Ferrari, D., and Rossi, F. (1995) *Nature* 374: 647-650.
45. Mattson, M. P., Cheng, B., Culwell, A. R., Esch, F. S., Lieberburg, I., and Rydel, R. E. (1993) *Neuron* 10: 243-254.
46. Furukawa, K., Sopher, B. L., Rydel, R. E., Begley, J. G., Pham, D. G., Martin, G. M., Fox, M., and Matteson, M. P. (1996) *Neurochem.* 67: 1882-1896.
47. Lannfelt, L., Basun, H., Wahlund, L.-O., Rowe, B. A., and Wagner, S. L. (1995) *Nature Med.* 1: 829-832.
48. Felsenstein, K.M., Hunihan, L.W. and Roberts, S.B. 1994. Altered cleavage and secretion of a recombinant β -APP bearing the Swedish familial Alzheimer's disease mutation. *Nature Genetics* 6: 251-256.

As will be recognized by those skilled in the art, the various embodiments described herein are provided by way of illustration and not limitation; various modifications and substitutions may be made thereto without departing from the spirit and scope of the invention. Such modifications and substitutions are contemplated as within the scope of the following claims.

WHAT IS CLAIMED IS:

I. A compound of FORMULA I, II or III:



or a pharmaceutically acceptable derivative, isostere, mixture or stereoisomer thereof, wherein:

R_1 comprises a W-substituted or unsubstituted lower alkenyl or a lower alkynyl, wherein said W-substituent comprises a halo, a hydroxy, an alkyl, an aryl, a heterocycle, a heteroaryl, an alkoxy, an aminosulfonyl, an alkylcarbonyl, an arylcarbonyl, a heteroarylcarbonyl, a heterocycle-carbonyl, a nitro, a haloalkyl, an amino or aminocarbonyl;

X comprises $-\text{CH} = \text{CH}-(\text{CH} = \text{CH})_k-(\text{CH}_2)_j$ or $-\text{CH}_2\text{CH}_2-(\text{CH}_2)_b$, wherein k comprises an integer selected from within the range of 0 to 9, j comprises an integer

selected from within the range of 0 to 4, *b* comprises an integer selected from within the range of 3 to 6;

Y comprises S, SO, SO₂, NR₂₀, -N(O)(R₂₀), -PR₂₀, -P(O)(R₂₀)-, -P(O)O- or -P(O)(OR₂₀)O, wherein R₂₀ comprises either a hydrogen atom or an alkyl or an aryl;

Z comprises -(CH₂)_{*i*}-A, wherein *i* is an integer selected from the range of 0 to 4 and A comprises an L- substituted or unsubstituted heterocycle, a heteroaryl or an aryl, and further, wherein said L-substituent comprises a halogen, a lower alkyl, a lower alkenyl, a lower alkynyl, an alkoxy, an aryloxy, a hydroxy, a haloalkyl, a trifluoromethyl, a cyano, a nitro, a nitrile, an alkylthio, a phenyl and an -NR₃₀R₃₁ group and wherein R₃₀ and R₃₁ each independently comprise H, alkyl, hydroxy or halo lower alkyl;

R₂, R₃, R₄, R₅, R₆, R₇, R₈, Q each independently comprise constituents, and *n* comprises zero or an integer, wherein each of said constituents is selected according to a group selected from among groups (i), (ii), (iii), (iv), (v), (vi) and (vii) as follows:

(i) R₃ and R₅ each independently comprise a side chain of a naturally occurring α-amino acid, a hydrogen atom, an alkyl, an alkenyl, an alkynyl, an aryl, an aralkyl, an aralkenyl, an aralkynyl, a heteroaryl, a heteroaralkyl, a heteroaralkenyl, a heteroaralkynyl, an L-substituted aryl, an L-substituted aralkyl, an L-substituted aralkenyl, an L-substituted aralkynyl, an M-substituted heteroaryl, an M-substituted heteroaralkyl, an M-substituted heteroaralkenyl or an M-substituted heteroaralkynyl, wherein M comprises a lower alkyl or a halo lower alkyl;

R₂, R₄, R₆, and R₈ each independently comprise a hydrogen atom, a lower alkyl, a cycloalkyl or a cycloalkylalkyl;

R₇ comprises a C₁₋₆ alkyl, an R_AR_BCH constituent, a aryl, an alkenyl, an alkynyl, a 9-fluorenyl, an aralkyl, an aralkenyl, an aralkynyl, a monocyclic or a bicyclic heterocycle or a monocyclic or a bicyclic aryl, heteroaryl or aryl-heteroaryl, wherein said aryl, said heterocycle and said heteroaryl groups are optionally substituted with L;

Q comprises a -C(O)-, a -O-C(O)-, an -S(O)₂- or a -HN-C(O)-;

n is zero or the integer 1;

R_A comprises a -(T)_{*m*}-(D)_{*m*}-R₃ constituent, wherein T is O or NH, D is C₁₋₄ alkyl or C₂₋₄ alkenyl, and *m* is zero or the integer 1; and

R_B comprises a constituent selected as set forth for R₃ and R₅; or alternatively,

(ii) R₂, R₅ and R₈ each independently comprise a constituent as set forth in (i);

R₃ and R₄ each independently comprise constituents as recited in (i) or (iv);

n is zero; and

R₆ and R₇ together with the atoms to which each is attached form a 3-10 membered heterocyclic moiety containing 1-5 heteroatoms; wherein said moiety is optionally substituted with L; or alternatively,

(iii) R₂, R₃, R₄, R₅ and R₈ each independently comprise a constituent as set forth in (i);

Q comprises a -C(O)-;

n comprises the integer 1; and

R₆ and R₇ each independently comprise a substituted or unsubstituted carbonyl (-(C=O)-), phenylene, heteroatom, lower alkylene, lower alkylene linked to a heteroatom, wherein said substituent comprises L and also wherein R₆ and R₇ together with the atoms to which they are attached form a cyclic or a bicyclic moiety; or alternatively,

(iv) R₂, R₅, R₆, R₇, R₈ and Q each independently comprise a constituent as set forth in any of (i)-(iii) or (v-vii), and n comprises zero or 1;

R₃ and R₄ each independently comprise a heteroatom or a substituted or unsubstituted lower alkylene or lower alkylene linked to a heteroatom, wherein said substituent comprises L and also wherein R₃ and R₄ together with the atoms to which they are attached form a heterocyclic moiety; or alternatively,

(v) R₂, R₇, R₈, Q each independently comprise a constituent as set forth in (i), and n comprises zero or 1;

R₃ and R₄ each independently comprise a constituent as set forth in either of (i) or (iv);

R₅ and R₆ each independently comprise a heteroatom or a substituted or unsubstituted lower alkylene or a lower alkylene linked to a heteroatom, wherein said substituent comprises L and also wherein R₅ and R₆ together with the atoms to which they are attached form a heterocyclic moiety; or alternatively,

(vi) R₂, R₆ and R₈ each independently comprise a constituent as set forth in (i);

R₃ and R₄ each independently comprise a constituent as set forth in either of (i) or (iv);

n is zero; and

R₅ and R₇ each independently comprise a heteroatom or a substituted or unsubstituted lower alkylene or lower alkylene linked to a heteroatom, wherein said substituent comprises L and also wherein R₅ and R₇ together with the atoms to which they are attached form a heterocyclic moiety; or alternatively,

(vii) R₂, R₅ and R₈ each independently comprise constituents as set forth in (i);

R₃ and R₄ each independently comprise constituents as set forth in either of (i) or (iv);

n is 0;

R₆ and R₇ each independently comprise a heteroatom or a substituted or unsubstituted lower alkylene or lower alkylene linked to a heteroatom, wherein said substituent comprises L and also wherein R₆ and R₇ together with the atoms to which each is attached form a 6-12 membered bicyclic heterocyclic or heteroaryl moiety.

2. The compound of claim 1, wherein:

R₂, R₃, R₄, R₅, R₆, R₇, R₈, Q wherein the constituents in said groups (i)-(vii) are further limited as follows;

(i) wherein when either of R₃ or R₅ comprise an alkyl, each independently comprise a lower alkyl; when R₃ or R₅ comprise an alkenyl, each independently comprises a C₂₋₁₀ alkenyl; when R₃ or R₅ comprise an alkynyl each independently comprises a C₂₋₆ alkynyl; wherein said M constituent comprises a C₁₋₄ alkyl or a trifluoromethyl; when R₂, R₄, R₆, or R₈ comprise an alkyl, each independently comprises a C₁₋₄ alkyl; when R₂, R₄, R₆, or R₈ comprise an aryl, each independently comprises an L-substituted aryl; when R₂, R₄, R₆, or R₈ comprise a heterocycle each independently comprises an L-substituted heterocycle; when R₂, R₄, R₆, or R₈ comprise a heteroaryl, each independently comprises an L-substituted heteroaryl;

(ii) wherein R₆ and R₇ each independently comprise a morpholino, a thiomorpholino, a pyrrolidinyl or a 4-hydroxypyrrolidinyl;

(iii) wherein R₆ and R₇ each independently comprise a succinimidyl, a phthalimidyl or a maleimidyl constituent;

(iv) R₃ and R₄ each independently comprise a morpholino, a thiomorpholino, a pyrrolidinyl or a 4-hydroxypyrrolidinyl; or alternatively,

(v) R_5 and R_6 each independently comprise a morpholino, a thiomorpholino, a pyrrolidinyl or a 4-hydroxypyrrolidinyl constituent;

(vi) R_5 and R_7 each independently comprise a morpholino, a thiomorpholino, a pyrrolidinyl or a 4-hydroxypyrrolidinyl; or alternatively,

(vii) R_6 and R_7 each independently comprise 1,2,3,4-tetrahydroisoquinolinyl.

3. The compound of claim 1, wherein k and j are 0 and b is 3 or 4.

4. The compound of claim 1, wherein X comprises $-\text{CH}=\text{CH}-$.

5. The compound of claim 1, wherein R_1 comprises a lower alkenyl.

6. The compound of claim 1, wherein Y comprises a SO , an SO_2 , an $-\text{N}(\text{O})(\text{R}_{20})$ constituent, a $-\text{P}(\text{O})(\text{R}_{20})$ - constituent or a $-\text{P}(\text{O})(\text{OR}_{20})$ - constituent.

7. The compound of claim 1, wherein Z comprises an L -substituted or unsubstituted phenyl, pyridyl, pyrazinyl, thienyl, furyl, pyrrolyl, imidazolyl or pyrimidinyl constituent.

8. The compound of claim 1, wherein Y comprises SO_2 .

9. The compound of claim 1, wherein X comprises $-\text{CH}=\text{CH}-$ and wherein, all constituents are as set forth in claim 1, but with the provisos that: Y does not comprise NR_{20} and A comprises an L -substituted or unsubstituted phenyl, pyridyl, pyrazinyl, thienyl, furyl, pyrrolyl, imidazolyl or pyrimidinyl, wherein said L does not comprise aryloxy.

10. The compound of claim 9, with the further provisos in said groups (i)-(vii) as follows:

(i) when R_3 or R_5 comprise an alkenyl or an alkynyl, each independently comprises a C_{2-10} alkenyl or a C_{2-6} alkynyl, and M comprises a C_{1-4} alkyl or a trifluoromethyl;

(ii) when R_2 , R_4 , R_6 , or R_8 comprise a lower alkyl, each independently comprises a C_{1-4} alkyl and when R_6 and R_7 together with the atoms to which each is attached form a heterocyclic moiety, said moiety comprises a morpholino, a thiomorpholino, pyrrolidinyl or a 4-hydroxypyrrolidinyl constituent;

(iii) when R_6 and R_7 together with the atoms to which they are attached form a cyclic or bicyclic moiety, said moiety comprises a succinimidyl, a phthalimidyl or a maleimidyl constituent;

(iv) when R_3 and R_4 together with the atoms to which they are attached form a heterocyclic moiety, said moiety comprises a morpholino, a thiomorpholino, pyrrolidinyl or a 4-hydroxypyrrolidinyl constituent;

(v) when R_5 and R_6 together with the atoms to which they are attached form a heterocyclic moiety, said moiety comprises a morpholino, a thiomorpholino, pyrrolidinyl or a 4-hydroxypyrrolidinyl constituent;

(vi) when R_5 and R_7 together with the atoms to which they are attached form a heterocyclic moiety, said moiety comprises a morpholino, a thiomorpholino, pyrrolidinyl or a 4-hydroxypyrrolidinyl constituent; and

(vii) when R_6 and R_7 together with the atoms to which they are attached form a heteroaryl moiety, said moiety comprises a 1,2,3,4-tetrahydroisoquinolinyl constituent.

11. The compound of claim 9, wherein: R_1 comprises an unsubstituted lower alkenyl or lower alkynyl and Y comprises SO or SO₂.

12. The compound of claim 11, wherein R_1 comprises an unsubstituted lower alkenyl and Y comprises SO₂.

13. The compound of claim 12, wherein: R_1 comprises 2-methylpropenyl or 2-butenyl; R_2 , R_4 , R_6 and R_8 each independently comprise a C₁₋₄ alkyl, a lower cycloalkyl or a lower cycloalkylalkyl; R_3 and R_5 each independently comprise a hydrogen atom, a C₁₋₄ alkyl, an alkylthio-alkyl, an aryl, an aralkyl, a heteroaralkyl, a 1-aminobut-4-yl or an aminocarbonyl-CH₂- constituent; R_7 and (Q)_n taken together comprise an acetyl, a benzyloxycarbonyl, an O-(3-chlorophenyl)-lactyl, a 9-fluorenylmethylcarbonyl, a Boc, 4-benzylpiperidinocarbonyl, a 4-benzylpiperazinocarbonyl or 1,2,3,4-tetrahydroisoquinolinylcarbonyl; and Z comprises an L-substituted or unsubstituted phenyl.

14. The compound of claim 9, wherein R_1 comprises 2-methylpropenyl.

15. The compound of claim 9, wherein: R_1 comprises 2-methylpropenyl; R_2 , R_4 , R_6 and R_8 each independently comprise a hydrogen atom, a cyclopropyl or a cyclopropyl-methyl; R_3 comprises isobutyl, benzyl, (2-methylthio) ethyl or isopropyl; R_5 comprises an isopropyl or an isobutyl; R_7 and (Q)_n taken together comprise benzyloxycarbonyl, 4-benzylpiperidinocarbonyl, 4-benzylpiperazinocarbonyl or 1,2,3,4-tetrahydro-isoquinolinylcarbonyl; Y comprises SO₂; and Z comprises a phenyl or a 4-(3-hydroxy-1-propynyl)phenyl.

16. The compound of claim 1 selected from the group consisting of: N-Cbz-L-Leu-L-Leu-L-Leuene phenyl vinyl sulfone; N-Cbz-L-Leu-L-Leu-D-Leuene phenyl vinyl sulfone; N-(THIQ-carbonyl)-L-Leu-L-Leu-D-Leuene phenyl vinyl sulfone; N-(THIQ-carbonyl)-L-Leu-L-Leu-L-Leuene phenyl vinyl sulfone; N-(THIQ-carbonyl)-L-Val-L-Met-L-Leuene phenyl vinyl sulfone; N-(THIQ-carbonyl)-L-Val-L-Met-D-Leuene phenyl vinyl sulfone; N-(THIQ-carbonyl)-L-Val-L-Leu-L-Leuene phenyl vinyl sulfone; N-(THIQ-carbonyl)-L-Val-L-Leu-D-Leuene phenyl vinyl sulfone; N-(4-benzylpiperidinylcarbonyl)-L-Leu-L-Leu-L-Leuene phenyl vinyl sulfone; N-(4-benzylpiperidinylcarbonyl)-L-Leu-L-Leu-D-Leuene phenyl vinyl sulfone; N-(4-benzylpiperazinylcarbonyl)-L-Leu-L-Leu-L-Leuene phenyl vinyl sulfone and N-(4-benzylpiperazinylcarbonyl)-L-Leu-L-Leu-D-Leuene phenyl vinyl sulfone.

17. A pharmaceutical composition comprising one or more compounds of claim 1, or a pharmaceutically acceptable derivative, isostere, mixture or stereoisomer thereof, in a pharmaceutically acceptable carrier.

18. A pharmaceutical composition comprising one or more compounds of claim 9, or a pharmaceutically acceptable derivative, isostere, mixture or stereoisomer thereof, in a pharmaceutically acceptable carrier.

19. A method of modulating processing of amyloid precursor protein to A β in an *in vitro* culture of cells, comprising the step of administering to said culture of cells the composition of claim 1.

20. A method of modulating processing of amyloid precursor protein to A β in an *in vitro* culture of cells, comprising the step of administering to said culture of cells the composition of claim 9.

21. A method of treating a subject in need thereof to alter processing of amyloid precursor protein to A β , comprising the step of administering to said subject the composition of claim 17.

22. A method of treating a subject in need thereof to alter processing of amyloid precursor protein to A β , comprising the step of administering to said subject the composition of claim 18.

23. The method of claim 21, wherein said subject in need thereof comprises a subject diagnosed as having a neurodegenerative disease selected from among: Alzheimer's disease, cognition deficits, Down's Syndrome, Parkinson's disease, cerebral hemorrhage with amyloidosis, dementia pugilistica and head trauma.

24. The method of claim 22, wherein said subject in need thereof comprises a subject diagnosed as having a neurodegenerative disease selected from among: Alzheimer's disease, cognition deficits, Down's Syndrome, Parkinson's disease, cerebral hemorrhage with amyloidosis, dementia pugilistica and head trauma.

25. The method of claim 21, wherein said subject in need thereof comprises a subject diagnosed as having an autoimmune disease selected from among: rheumatoid arthritis, systemic lupus erythematosus, Sjogren's syndrome, insulin dependent diabetes type I or type II,

26. The method of claim 22, wherein said subject in need thereof comprises a subject diagnosed as having an autoimmune disease selected from among: rheumatoid arthritis, systemic lupus erythematosus, Sjogren's syndrome, insulin dependent diabetes type I or type II,

27. The method of claim 21, wherein said subject in need thereof comprises a subject diagnosed as having a traumatic tissue injury, a cardiovascular insufficiency, a stroke, a glucose tissue deprivation syndrome.

28. The method of claim 22, wherein said subject in need thereof comprises a subject diagnosed as having a traumatic tissue injury, , a cardiovascular insufficiency, a stroke, a glucose tissue deprivation syndrome.

29. A method for inhibiting the activity of a cysteinyl protease, comprising the step of bringing the cysteinyl protease into contact with a compound of claim 1.

30. A method for inhibiting the activity of a cysteinyl protease, comprising the step of bringing the cysteinyl protease into contact with a compound of claim 9.

31. An article of manufacture, comprising packaging material and a compound of claim 1, contained within the packaging material, wherein said compound is effective to ameliorate or prevent one or more symptoms of a neurological disease, and the packaging material includes a label and a package insert that indicates that the compound is useful for treatment or prevention of said neurological disorder. .

32. An article of manufacture, comprising packaging material and a compound of claim 9, contained within the packaging material, wherein said compound is effective to ameliorate or prevent one or more symptoms of a neurological disease, and the packaging material includes a label and a package insert that indicates that the compound is useful for treatment or prevention of said neurological disorder.

INTERNATIONAL SEARCH REPORT

International application no.

PCT/US00/16864

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07K 5/08

US CL : 530/331

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/331; 514/18, 19

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST, CAS Online

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,856,530 A (WEBBER et al) 05 January 1999, see entire document.	1-32

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
C document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

28 AUGUST 2000

Date of mailing of the international search report

03 OCT 2000

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DAVID LUKTON

Telephone No. (703) 308-0196